

GENETIC TOLERANCE TO FERROUS AND ALUMINIUM TOXICITIES FOR
SEED GERMINATION IN *Oryza sativa* L.

NUSRAT JAHAN

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Doctor of Philosophy (Bioscience)

Faculty of Biosciences and Medical Engineering
Universiti Teknologi Malaysia

MAY 2017

Specially dedicated to my Mother (*Mor*) & Father (*Plar*)

ACKNOWLEDGEMENT

In the name of ALLAH (S.W.T), the most Gracious and the most Compassionate. This PhD or anything without ALLAH's help is impossible. I thank you for ALLAH (S.W.T) help and I would always like to be your obedient servant.

This PhD research study would not have been possible without the cooperation and support by my supervisor Dr. Muhammad Arshad Javed. His patience, during the numerous focus group discussions as well the lab work help to understand things in more easy way. I am sincerely apologizing for any of my actions that made him bother during my research.

Sincerely thanks to my co-supervisors: Dr. Azman Abd Samad and Dr. Fazilah Bt. Abd Manan for their kind support and guidance throughout my PhD.

A very special thank you to Dr. Rashid Ahmed (Department of Physics, Faculty of Science, Universiti Teknologi Malaysia) for his invaluable support to allowing me to compute and analyze my data results in Scientific Computing and Instrumentation Lab.

My deep appreciation goes to my research fellows and colleagues: Samiullah Khan and Muhammad Waseem Chughtai, as well as local research fellows: Ms.Farah and Mrs.Atiqah for their support and companionship.

Lastly, I would like to thank my all family group for their love and encouragement. For my parents, who raised me with a love of science and supported me in all my pursuits. And most of all for my loving, supportive, encouraging, and patient brothers, beloved sister, aunts, and uncles (especially Nadir Shah) who's faithful support during this difficult period of my Ph.D. Thank you!

ABSTRACT

Direct seeded culture is famous for rice cultivation due to its low inputs and less labour demand. However seeds are exposed to various abiotic stresses in direct seeding culture. Ferrous (Fe^{2+}) and aluminium (Al^{3+}) are toxic metals that severely affect seed germination and growth traits. Genetic tolerance to metal toxicity would be one of the possible solutions to combat with these challenges. The genetic study on Fe^{2+} and Al^{3+} for seed germination can hardly be found in literature. The present study was conducted to identify the potential loci linked with genes controlling tolerance to Fe^{2+} and Al^{3+} for seed germination. A segregating mapping population of $\text{F}_{2:3}$ was developed from the cross between selected Fe^{2+} and Al^{3+} tolerant parent *Pokkali* and sensitive parent, *Pak Basmati*. Parental DNA was used for polymorphic survey and F_2 DNA was used for genotyping. A molecular linkage map was constructed using 84 markers data. The molecular linkage map covered 3435.5cM with an average distance of 7.63cM except 4 larger gaps on chromosome 1, 2, and 4. F_3 progenies (129) were evaluated against the optimized toxic level of Fe^{2+} and Al^{3+} under controlled environment. Germination traits such as final germination percentage (FG%), germination energy (GE), germination rate (GR), germination speed (GS), germination index (GI), mean time of germination (MGT), germination value (GV), germination velocity (GVe), peak value of germination (PV), germination capacity (GC) and growth traits such as root length (RL), shoot length (SL), total dry biomass (TDB), and germination vigour index (GVI) were measured. In the present study, screening of six rice varieties showed significant differences for seed germination, however *Pokkali* exhibited the minimum and *Pak Basmati* showed the maximum influence in seed germination. A 20mM (at pH4.0) of both Fe^{2+} and Al^{3+} was found to be the optimized toxic level, as most germination and growth parameters were found to significantly affected at this concentration. Phenotypic data showed significant variations in germination and growth parameters. Total of 39 QTLs (Quantitative Trait Loci) for germination traits and 8 QTLs for growth parameters linked with Fe^{2+} toxicity tolerance were determined by Simple Interval Mapping (SIM) at 0.06% to 39.9% of phenotypic variations, respectively. Thirty four QTLs for germination parameters and 8 QTLs for growth traits with phenotypic variations 0 to 47% traits linked with Fe^{2+} toxicity tolerance were mapped by Composite Interval Mapping (CIM). Forty nine QTLs for germination and 23 putative QTLs for growth parameters with phenotypic variations 0.01% to 69% were determined using Multiple Interval Mapping (MIM). Epistasis analysis revealed that the QTLs are mostly dependent on the alleles at other loci. For Al^{3+} toxicity tolerance 40 markers linked with germination and growth parameters with phenotypic variation 0 to 62.64% were identified by Simple Interval Mapping. Forty-six putative QTLs with phenotypic variation 0 to 28.1% were detected for germination and growth parameters for Al^{3+} toxicity tolerance using Composite Interval Mapping. Sixty five markers linked with germination and growth traits, at 0 to 72.7% phenotypic variations were mapped by Multiple Interval Mapping. Epistasis analysis showed that Al^{3+} toxicity tolerance is polygenic and controlled by additive effect. The results suggested that the QTL regions (18cM) and (72cM) were tightly linked to Al^{3+} tolerance genes could be used for marker assisted selection programme using fine mapping. Moreover, this study also provides an understanding to exclusion tolerance mechanism of Al^{3+} toxicity as known in the *Triticeae* within sub-family *Pooideae*. The identified major QTLs of this research would be useful for rice hybridization programs to induce tolerance against these toxic metals.

ABSTRAK

Budaya pembenihan secara terus sangat terkenal dalam penanaman padi kerana input yang rendah dan permintaan buruh yang kurang. Walau bagaimanapun benih terdedah kepada pelbagai tekanan abiotik dalam budaya tabur terus. Ferus (Fe^{2+}) dan aluminium (Al^{3+}) adalah logam toksik yang mempengaruhi percambahan benih dan pertumbuhan sifat-sifatnya. Genetik ketoksikan logam merupakan salah satu penyelesaian untuk mengatasi cabaran ini. Kajian genetik pada Fe^{2+} dan Al^{3+} untuk percambahan benih belum banyak dikaji dalam kajian lepas. Kajian ini dijalankan untuk mengenal pasti potensi lokus yang dikaitkan dengan toleransi kawalan gen Fe^{2+} dan Al^{3+} dalam proses percambahan benih. Pemetaan pengasingan kelompok $\text{F}_{2:3}$ telah dibangunkan daripada penyilangan antara Fe^{2+} terpilih dan *Pokkali* toleran dasar Al^{3+} serta *Pak Basmati*. DNA asal telah digunakan untuk kajian polimorfik dan DNA F_2 telah digunakan untuk penjenisan gen. Peta Rangkaian molekul telah dibina menggunakan 84 data penanda. Peta hubungan molekul merangkumi 3435.5cM dengan jarak purata 7.63cM kecuali 4 jurang lebih besar pada kromosom satu, dua dan empat. Progeni F_3 (129) telah dinilai berdasarkan tahap toksik yang dioptimumkan Fe^{2+} dan Al^{3+} di bawah kawalan persekitaran. Sifat-sifat percambahan seperti peratusan percambahan akhir (FG%), percambahan tenaga (GE), kadar percambahan (GR), kelajuan percambahan (GS), indeks percambahan (GI), min masa percambahan (MGT), nilai percambahan (GV), halaju percambahan (G_{Ve}), nilai puncak percambahan (PV), kapasiti percambahan (GC) dan pertumbuhan ciri-ciri seperti tempoh asal (RL), tempoh bertunas (SL), jumlah bahan kering (TDB), dan indeks percambahan tenaga (GVI) telah diukur. Dalam kajian ini, enam variasi padi menunjukkan perbezaan yang signifikan bagi percambahan benih, bagaimanapun *Pokkali* dipamerkan secara minimum dan *Pak Basmati* menunjukkan pengaruh maksimum dalam percambahan benih. Sebanyak 20mM (pada pH4.0) tahap toksik pada kedua-dua Fe^{2+} dan Al^{3+} ditemui secara optimum kerana kebanyakan percambahan dan pertumbuhan parameter didapati terjejas dengan ketara pada kepekatan ini. Data fenotip menunjukkan perbezaan ketara dalam percambahan dan pertumbuhan parameter. Sebanyak tiga puluh sembilan QTLs (Ciri Kuantatif Loci) untuk ciri-ciri percambahan dan lapan QTLs untuk parameter pertumbuhan dikaitkan dengan Fe^{2+} ketoksikan toleransi telah ditentukan oleh selang ringkas pada 0.06% kepada 39.9% daripada variasi fenotip. Tiga puluh empat QTLs parameter percambahan dan lapan QTLs ciri-ciri pertumbuhan dengan variasi fenotip 0-47% ciri-ciri yang berkaitan dengan Fe^{2+} ketoksikan toleransi telah dipetakan oleh Selang Komposit Pemetaan (CIM). Empat puluh sembilan QTLs percambahan dan dua puluh tiga QTLs untuk parameter pertumbuhan variasi fenotip 0.01%-69% telah ditentukan menggunakan Selang Pemetaan Pelbagai (MIM). Analisis epistasis menjelaskan bahawa kebanyakan QTLs bergantung kepada alel pada lokus yang lain. Untuk ketoksikan Al^{3+} empat puluh toleransi penanda dikaitkan dengan percambahan dan pertumbuhan parameter dengan fenotip perubahan 0-62.64% telah dikenal pasti oleh Selang Pemetaan Mudah (SIM). Empat puluh enam QTLs dengan fenotip Perubahan 0-28.1% dikesan untuk percambahan dan pertumbuhan parameter bagi Al^{3+} ketoksikan toleransi menggunakan Komposit Selang Pemetaan. Enam puluh lima penanda dikaitkan dengan percambahan dan pertumbuhan ciri-ciri iaitu 0-72.7% variasi fenotip telah dipetakan oleh Selang Pemetaan Pelbagai. Analisis epistasis menunjukkan bahawa Al^{3+} ketoksikan toleransi adalah poligenik dan dikawal oleh kesan tambahan. Dapatan kajian mencadangkan bahawa kawasan QTL (18cm) dan (72cm) yang berkait rapat dengan Al^{3+} gen toleransi boleh digunakan sebagai penanda program pemilihan yang dibantu dengan menggunakan pemetaan halus. Selain itu, kajian ini juga memberi kefahaman kepada mekanisme pengecualian toleransi Al^{3+} ketoksikan yang diketahui dalam *Triticeae* sub-keluarga padi *Pooideae*. QTLs utama yang dikenal pasti dalam kajian ini akan digunakan untuk program penghibridan beras bagi mendorong toleransi terhadap logam toksik ini.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENT	vii
	LIST OF TABLES	xiv
	LIST OF FIGURES	xvii
	LIST OF ABBREVIATION	xxi
	LIST OF APPENDICES	xxiv
1	INTRODOCUTION	1
	1.1. Research Background	1
	1.2. Problem Statement	4
	1.3. Research Objectives	5
	1.4. Scope of Study	6
	1.5. Significance of Study	6
2	LITERATURE REVIEW	7
	2.1. Overview	7
	2.2. Toxic Soils	8

2.2.1.	Acid Soil	9
2.3.	Iron (Fe^{2+}) Toxicity	11
2.4.	Aluminum (Al^{3+}) Toxicity	13
2.5.	Plant Tolerant / Resistance Mechanisms for Fe^{2+} and Al^{3+}	14
2.5.1.	Iron (Fe^{2+}) Tolerance / Resistant Mechanisms	14
2.5.1.1.	Oxidation of Fe^{2+} at Root Surface	15
2.5.1.2.	Exclusion of Fe^{2+} at Root Surface	16
2.5.1.3.	Retention of Fe^{2+} in the Root Tissue	16
2.5.1.4.	Retention of Fe^{2+} in Apoplast of Leaf	17
2.5.1.5.	Symplastic Tissue Tolerance Against Fe^{2+} Toxicity	17
2.5.2.	Al^{3+} Tolerance / Resistance Mechanisms	17
2.5.2.1.	Exclusion of Al^{3+} Outside the Root	18
2.5.2.2.	Symplastic Tolerance (Al^{3+} Accumulation in the Symplasm)	19
2.5.3.	Seed Germination Tolerance / Resistant Mechanisms	20
2.6.	Tolerant Genes in Crop Plants	21
2.7.	Germination Parameters to Characterize the Seed Toxicity Effects	22
2.8.	Molecular Markers and Linkage Maps	25
2.9.	Mapping Populations	26
2.10.	MAPMAKER/EXP	27

2.11.	Quantitative Trait loci (QTL) Mapping: Methodology and Application	27
2.12.	Applications of QTL Analysis	29
2.12.2.	Marker Assisted Selection (MAS)	29
2.12.3.	QTL for Fine Mapping	30
2.12.4.	The Combination of QTL Mapping Transcriptome Profiling	30
2.12.5.	Pyramiding	31
2.13.	QTL Analysis for Tolerance to Fe^{2+} and Al^{3+} Toxicity in Rice	31
2.14.	The Genetics and Molecular Studies of Fe^{2+} and Al^{3+} Tolerance	33
3	RESEARCH METHODOLOGY	35
3.1.	Overview	35
3.2.	Construction of Mapping Population	35
3.2.1.	Selection of Rice Genotypes for Parental Line Under Fe^{2+} and Al^{3+} Toxicities	36
3.2.2.	Hybridization of Selection Parent and Molecular Pedigree Evaluation	36
3.2.2.1.	DNA Extraction	37
3.2.2.2.	DNA Quality Analysis by Agarose Gel	38
3.2.2.3.	Estimation of Quantity, Quality, and Purity of DNA	38
3.2.2.4.	Screening of Polymorphisms Among Parental Line (<i>Pak Basmati</i> and <i>Pokkali</i>)	39
3.2.2.5.	PCR (DNA amplification) PCR Amplification	39

3.2.2.6. Gel Electrophoresis	40
3.2.2.7. Electrophoresis	40
3.2.2.8. Silver Staining	41
3.2.2.9. Scoring of Bands	42
3.2.3. Mapping Population	42
3.3. Tolerance Limits Optimization for Parental Line Against Fe^{2+} and Al^{3+} Toxicities	43
3.3.1. Maintaining pH of Iron Solutions	44
3.3.2 Phenotyping of F_3 Progenies	45
3.4. Construction of Linkage Map	46
3.4.1. Genotyping of F_2 Population with SSR	47
3.4.1.1. Screening of Polymorphism Among Parental Line (<i>Pak Basmati</i> and <i>Pokkali</i>)	47
3.4.1.2. PCR (DNA) Amplification	47
3.4.1.3. Gel Electrophoresis	48
3.4.1.4. Scoring of Bands	48
3.4.1.5. Segregation Distortion	48
3.4.2. Construction of Molecular Linkage Map	49
3.5. Identification of Microsatellite Loci Linked to Tolerance to Fe^{2+} and Al^{3+} Toxicities	49
3.6. QTL Nomenclature	50
3.7. Statistical Analysis	50
4 RESULTS AND DISCUSSION	53
4.1. Iron (Fe^{2+}) Toxicity	53

4.1.1. Screening of Rice Genotypes for Parental Line Selection in Fe^{2+} Toxicity	53
4.1.2. Selection of Parental Line for Mapping Population	60
4.1.3. Hybridization of Selected Parent and Molecular Pedigree Evaluation	60
4.1.3.1. F_1 Hybrid Confirmation	62
4.1.3.2. Construction of Mapping Population	63
4.1.4. Optimization of Tolerance Limit Against Fe^{2+} Toxicity in Parental Line and to Phenotype the F_2 Against Optimized Level	64
4.1.4.1. ANOVA for Germination and Growth Parameters at 5mM and 10mM of FeCl_2	64
4.1.4.2. ANOVA for Germination and Growth Parameters at 15mM and 20mM of FeCl_2	65
4.1.5. F_3 Phenotyping	70
4.1.5.1. Phenotypic Variation and Frequency Distribution	71
4.1.5.2. Correlation between all Germination and Seedling Growth Parameters	73
4.1.6. Construction of Molecular Linkage Map	77
4.1.6.1. DNA Extraction	77
4.1.6.2. Parental Polymorphic Survey of Parental Line	77
4.1.6.3. Marker Segregation	79
4.1.6.4. Linkage Map	81
4.1.7. QTL Analysis	84

4.1.7.1. Identification of Molecular Markers Linked to QTLs by Simple Interval Mapping (SIM)	84
4.1.7.2. Identification of Molecular Markers Linked to QTLs by Composite Interval Mapping (CIM)	95
4.1.7.3. Identification of Molecular Markers Linked to QTLs by Multiple Interval Mapping (MIM)	108
4.1.8. Summary for Identified QTLs Associated with Fe ²⁺ Toxicity Tolerance	129
4.2. Aluminum (Al ³⁺) Toxicity	130
4.2.1. Screening of Rice Genotypes for Parental Line Selection in Al ³⁺ Toxicity	130
4.2.2. Hybridization of Selected Parent and molecular Pedigree Evaluation	137
4.2.3. Optimization of Tolerance Limit Against Al ³⁺ Toxicity in Parental Line	137
4.2.3.1. ANOVA for Germination Parameters at 5mM and 10mM for AlCl ₃	137
4.2.3.2. ANOVA for Germination and Growth Parameters at 15mM and 20mM for AlCl ₃	140
4.2.4. F ₃ Phenotyping	143
4.2.4.1. Phenotypic Variation and Frequency Distribution	143
4.2.4.2. Correlation between all Germination and Seedling Growth Parameters	146
4.2.5. Construction of Molecular Linkage Map	150
4.2.6. QTL Analysis	150

4.2.6.1. Identification of Molecular Markers Linked to QTLs by Simple Interval Mapping (SIM)	150
4.2.6.2. Identification of Molecular Markers Linked to QTLs by Composite Interval Mapping (CIM)	163
4.2.6.3. Identification of Molecular Markers Linked to QTLs by Multiple Interval Mapping (MIM)	178
4.2.7. Summary for Identified QTLs Associated with Al ³⁺ Toxicity Tolerance	196
5 CONCLUSION AND FUTURE WORK	197
5.1. Future Work	198
REFERENCES	199
Appendices (A-H)	230-274

LIST OF TABLES

TABLE NO	TITLE	PAGE
2.1	Toxic elements concentration increase in low pH of soil	12
2.2	Abiotic stress-tolerant genes in crop plants (Ansari <i>et al.</i> , 2015)	22
3.1	Details of SSR markers used for F ₁ hybrid confirmation	39
3.2	Formulation of germination and growth parameters with references	45
4.1	Analysis of variance showing mean sum of squares of germination parameters for six rice varieties in different concentrations (0,3mM, 6mM, 9mM, and 12mM at pH 4.0) of Fe ²⁺ toxicity	54
4.2	Comparison of mean results for effect of various concentrations (0,3mM, 6mM, 9mM, and 12mM at pH 4.0) of Fe ²⁺ toxicity on germination parameters of six rice varieties	55
4.3	Coefficient of linear correlation between germination parameters of rice varieties at different concentrations (0,3mM, 6mM, 9mM, and 12mM at pH 4.0) of Fe ²⁺ toxicity	58
4.4	Analysis of variance of 5mM and 10mM of Fe ²⁺ toxicity at pH4.0 in germination parameters on parental line <i>Pak Basmati</i> and <i>Pokkali</i>	65
4.5	Analysis of variance of 15mM and 20mM of Fe ²⁺ toxicity at pH4.0 in germination parameters on parental line <i>Pak Basmati</i> and <i>Pokkali</i>	68

4.6	Coefficient of correlation between all germination and seedling growth parameters of F ₃ progenies in 20mM at pH 4.0 of Fe ²⁺ toxicity	75
4.7	Polymorphism survey for selection of appropriate SSR markers	79
4.8	Mapped markers distribution and distorted segregation	80
4.9	(a): QTLs identification for germination parameters linked to Fe ²⁺ toxicity tolerance using Simple Interval Mapping (SIM)	85
	(b): QTLs identification for growth parameters linked to Fe ²⁺ toxicity using Simple Interval Mapping (SIM)	90
4.10	(a): Composite Interval Mapping (CIM) analysis for germination parameter	96
	(b): Composite Interval Mapping (CIM) analysis for growth parameters	101
4.11	(a): Multiple Interval Mapping (MIM) for germination parameters	109
	(b): Multiple Interval mapping (MIM) analysis for growth parameters	118
4.12	Summary of total identified QTLs liked with tolerance to Fe ²⁺ toxicity by SIM, CIM, and MIM	129
4.13	Analysis of variance showing mean sum of squares of germination parameters for six rice varieties in different concentrations (0, 3mM, 6mM, 9mM, and 12mM) at pH 4.0 of Al ³⁺ toxicity	131
4.14	Comparison of mean for effect of various concentrations (0, 3mM, 6mM, 9mM, and 12mM) at pH4.0 Al ³⁺ toxicity at pH 4.0 on germination traits of six rice varieties	132
4.15	Coefficient of correlation between germination parameters of rice genotypes under Al ³⁺ toxicity	135

4.16	Analysis of variance of 5mM and 10mM of Al^{3+} toxicity in germination and growth parameters on parental line <i>Pak Basmati</i> and <i>Pokkali</i>	138
4.17	Analysis of variance of 15mM and 20mM of Al^{3+} toxicity in germination and growth parameters on parental line <i>Pak Basmati</i> and <i>Pokkali</i>	140
4.18	Coefficient of correlation between all germination and seedling growth parameters under 20mM Al^{3+} toxicity at pH4.0	147
4.19	(a): Simple Interval Mapping (SIM) analysis for germination parameter	151
	(b): QTLs identification for growth parameters linked to Al^{3+} toxicity using Simple Interval Mapping (SIM)	157
4.20	(a): Composite Interval Mapping (CIM) analysis for germination parameters	164
	(b): Composite Interval Mapping (CIM) analysis for growth parameters in Al^{3+} toxicity	170
4.21	(a): Multiple Interval Mapping (MIM) analysis and Epistasis for germination parameters under Al^{3+} toxicity	177
	(b): QTL analysis using Multiple Interval Mapping (MIM) and epistasis in Al^{3+} toxicity	185
4.22	Summary of total identified QTLs linked with tolerance to Al^{3+} toxicity by SIM, CIM, and MIM	196

LIST OF FIGURES

FIGURE NO	TITLE	PAGE
1.1	Problem statement diagram	4
3.1	Schematic representation: The progress of constructing mapping population	43
3.2	Research Operational Framework	52
4.1	Reduction percentage for germination parameters in six rice varieties for selection of parental lines	57
4.2	Hybridization of (a) <i>Pak Basmati</i> as Female and (b) <i>Pokkali</i> as Male and (c) the production of visible six F ₁ seeds for the development of mapping population	61
4.3	Polymorphic microsatellite marker profiles confirming the F ₁ hybridity. PB= <i>Pak Basmati</i> , Pok= <i>Pokkali</i> , F ₁ =Pok × PB individuals, (1, 2, 3, 4 defines F ₁₋₁ , F ₁₋₂ , F ₁₋₃ and F ₁₋₄), H=hybrid; S=self-pollinated, W=weight marker (100bp), bp=base pairs.	62
4.4	Seeds of F ₃ progenies	64
4.5	Effects of 5mM and 10mM Fe ²⁺ toxicity on parental line PB = <i>Pak Basmati</i> and Pok = <i>Pokkali</i>	66
4.6	Effects of 15mM and 20mM Fe ²⁺ toxicity on parental line PB = <i>Pak Basmati</i> and Pok = <i>Pokkali</i>	69
4.7	F ₃ phenotyping under optimized level of Fe ²⁺ toxicity, (a) P36 = number of F ₃ plants and (b) P37 = number of F ₃ plants	71
4.8	(a): Phenotypic variations and frequency distribution of Fe ²⁺ toxicity among 129 F _{2:3} for germination parameters. The arrows in black colour show mean of parental line. Pok= <i>Pokkali</i> , PB= <i>Pak Basmati</i>	72

	(b): Phenotypic variations and frequency distribution of Fe^{2+} toxicity among 129 $F_{2:3}$ for growth parameters. The arrows in black colour show mean of parental line. Pok= <i>Pokkali</i> , PB= <i>Pak Basmati</i>	73
4.9	Screening to identify the polymorphic loci between parental line, on chromosome 2 (rice genetic maps and genome sequence maps). 1= <i>Pokkali</i> , 2= <i>Pak Basmati</i> , W=weight marker 100bp, boxes shows polymorphism among parents	77
4.10	Silver-stained polyacrylamide gel electrophoresis showing the segregation patterns of F_2 population. Pok= <i>Pokkali</i> , PB= <i>Pak Basmati</i> , M=weight marker. L1,L2,L3=number of gel loadings	78
4.11	A molecular linkage map with 84 SSR marker loci constructed from 129 $F_{2:3}$ population line of the cross <i>Pak Basmati</i> \times <i>Pokkali</i> . The distances between markers are in centiMorgan (cM) Kosambi function using MAPMAKER/EXP. Left side of the map of linkage group are distances in cM among markers and on right side showing name of markers.	82
4.12	Fe^{2+} Chromosomal locations of QTLs by SIM analysis for FG%=Final germination percentage, GVe=germination velocity, GE=Germination energy, SG=Speed of germination, GPV=Germination peak value, GI= Germination index, GC=Germination capacity, GV=Germination value, MGT=Mean germination time, RL=Root length, SL=Shoot length, TDB=Total dry biomass, GVI=Germination Vigour index	93
4.13	Fe^{2+} Chromosomal locations of QTLs by CIM analysis for FG%=Final germination percentage, GVe=germination velocity, GE=Germination energy, SG=Speed of germination, GPV=Germination peak value, GI= Germination index, GC=Germination capacity, GV=Germination value, MGT=Mean germination time, RL=Root length, SL=Shoot length, TDB=Total dry biomass, GVI=Germination Vigour index	105
4.14	Fe^{2+} Chromosomal locations of QTLs by MIM analysis for FG%=Final germination percentage, GVe=germination velocity, GE=Germination energy, SG=Speed of germination, GPV=Germination peak value, GI= Germination index, GC=Germination capacity, GV=Germination value, MGT=Mean germination time, RL=Root length, SL=Shoot length, TDB=Total dry biomass, GVI=Germination Vigour index	124

4.15	Reduction percentage for germination parameters from (i) to (viii) in six rice varieties for selection of parental lines under Al^{3+} toxicity	134
4.16	Effects of 5mM and 10mM Al^{3+} toxicity on root length and shoot length of parental line <i>Pak Basmati</i> and <i>Pokkali</i>	139
4.17	Effects of 15mM and 20mM Al^{3+} toxicity parental line of <i>Pak Basmati</i> and <i>Pokkali</i>	141
4.18	F_3 phenotyping under optimized level of Al^{3+} toxicity, (a) P151 = number of F_3 plants and (b) P77 = number of F_3 plants	143
4.19	(a): Phenotypic variations and frequency distribution of Al^{3+} toxicity among 129 $F_{2:3}$ for germination parameters. The arrows in black colour show mean of parental line. Pok= <i>Pokkali</i> , PB= <i>Pak Basmati</i>	144
	(b): Phenotypic variations and frequency distribution of Al^{3+} toxicity among 129 $F_{2:3}$ for growth parameters. The arrows in black colour show mean of parental line. Pok= <i>Pokkali</i> , PB= <i>Pak Basmati</i>	145
4.20	F_3 plant with long root and well growth under Al^{3+} toxicity stress	146
4.21	Chromosomal locations of QTLs by SIM analysis linked to Al^{3+} toxicity tolerance for FG%=Final germination percentage, GVe=germination velocity, GE= Germination energy, SG=Speed of germination, GPV=Germination peak value, GI=Germination index, GC= Germination capacity, GV=Germination value, MGT=Mean germination time, RL=Root length, SL=Shoot length, TDB=Total dry biomass, VI=Vigour index	160
4.22	Chromosomal locations of QTLs by CIM analysis linked to Al^{3+} toxicity tolerance for FG%=Final germination percentage, GVe=germination velocity, GE=Germination energy, SG=Speed of germination, GPV=Germination peak value, GI=Germination index, GC=Germination capacity, GV=Germination value, MGT=Mean germination time, RL=Root length, SL=Shoot length, TDB= Total dry biomass, GVI=Germination Vigour index	173

- 4.23 Chromosomal locations of QTLs by MIM analysis linked to Al^{3+} toxicity tolerance for FG%=Final germination percentage, GVe=germination velocity, GE=Germination energy, SG=Speed of germination, GPV=Germination peak value, GI=Germination index, GC=Germination capacity, GV=Germination value, MGT=Mean germination time, RL=Root length, SL=Shoot length, TDB=Total dry biomass, GVI=Germination Vigour index

LIST OF ABBREVIATIONS

AFLPs	-	Amplified Fragment Length Polymorphisms
ANOVA	-	Analysis Of Variance
BC	-	Backcross
bp	-	Base pair
Chr	-	Chromosome
SIM	-	Simple Interval Mapping
CIM	-	Composite Interval Mapping
MIM	-	Multiple Interval Mapping
cM	-	centiMorgan
CTAB	-	Cetyl Trimethyl Ammonium Bromide
DH	-	Double Haploids
F _{2:3}	-	Second filial generation advanced to third filial generation
IRRI	-	International Rice Research Institute
RILs	-	Recombinant Inbred Lines
Pok	-	Pokkali
PB	-	Pak Basmati
BC1	-	First Backcross Generation
BC1F2	-	Second Backcross Generation after selfing

DH	-	Double haploid
MAS	-	Marker Assisted Selection
RFLP	-	Restriction Fragment Length Polymorphism
AFLP	-	Amplified Fragment Length Polymorphism
SSRs	-	Simple Sequence Repeats
SNPs	-	Single Nucleotide Polymorphisms
C	-	Codominant
D	-	Dominant
WinQTLCart	-	Windows Quantitative Trait Loci Cartographer
MapQTL	-	Mapping Quantitative Trait Loci
DHLs	-	Double Haploid Lines
LOD	-	Logarithm of Odd
IRRI	-	International Rice Research Institute
FG%	-	Final Germination Percentage
GE	-	Germination Energy
GVe	-	Germination Velocity
SG	-	Speed of Germination
GPV	-	Germination Peak Value
GC	-	Germination Capacity
GI	-	Germination Index
GV	-	Germination Value
MGT	-	Mean Germination Time
GVI	-	Germination Vigour Index

RL	-	Root Length
SL	-	Shoot Length
TDB	-	Total Dry Bio Mass
<i>r</i>	-	Pearson coefficient
Tris-HCl	-	Tris-Hydro Chloride
EDTA	-	Ethylenediaminetetraacetic acid
APS	-	Ammonium Persulfate
PIC	-	Polymorphism Information Content
χ^2	-	Chi Square
IM	-	Interval Mapping
R^2	-	Phenotypic Variation
LRT	-	Likelihood Ratio Test
RM	-	Rice Marker
Chr No	-	Chromosome Number

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Population line of F ₂ and F ₃ in Terms of the Philosophical Analysis	230
B	Required materials, chemical and solution preparation method	231
C	DNA Quantification Of F ₂ Population	234
D	List of 84 Polymorphic Markers	238
E	i. 5mM FeCl ₂ ANOVA Tests for Germination and Growth Parameters	243
	ii. 10mM FeCl ₂ ANOVA Tests for Germination and Growth Parameters	246
	iii. 15mM FeCl ₂ ANOVA Tests for Germination and Growth Parameters	250
	iv. 20mM FeCl ₂ ANOVA Tests for Germination and Growth Parameters	253
F	i. 5mM AlCl ₃ ANOVA Tests for Germination and Growth Parameters	257
	ii. 10mM AlCl ₃ ANOVA Tests for Germination and Growth Parameters	260
	iii. 15mM AlCl ₃ ANOVA Tests for Germination and Growth Parameters	263
	iv. 20mM AlCl ₃ ANOVA Tests for Germination and Growth Parameters	267
G	Research work pictures (outdoor/indoor)	271
H	List of published/submitted papers	274

CHAPTER 1

INTRODUCTION

1.1 Research background

Rice is one of the oldest crop plants cultivated since ancient time. It belongs to the genus *Oryza* of grass family (Gramineae). The genus *Oryza* consists of 21 wild and two cultivated species *Oryza sativa* and *Oryza glaberrima*. *Oryza sativa* is widely cultivated in Asia, America, and Europe. There are two subspecies of *Oryza sativa*, i.e. *Indica*, and *japonica* that comprised most varieties of rice and based (Oka, 1988). *Indica* and *Japonica* are further classified into temperate (*Japonica*), tropical (*Japonica*), indica (*Indica*) and *Aus* (*Indica*).

Rice (*Oryza sativa* L.) is a very important staple food for almost half of the world's population. Of the 49% of energy consumed by the world population in the form of cereal crops like rice, wheat, and corn, 23% is provided by rice. More than 90% rice is consumed and grown in Asia. It is grown in a large range of agricultural conditions between 55N and 35S altitude from sea level to 25000m or higher altitude (Subudhi *et al.*, 2006). Rice is cultivated on 11% of agricultural land in the world or about 154 million hectares, with 700 million tons of production annually (Halwart and Gupta, 2004). The tropical and subtropical climates are suitable for *indica* species, while *Japonica* cultivars are adapted to subtropical and temperate climates. Rice can be cultivated in four main ecosystems: a) upland; b) rainfed lowland; c) irrigated and d) flood prone. The soil texture, water holding capacity, and nutrient status are different in all rice ecosystems and is largely aerobic (Halwart and Gupta, 2004; IRRI, 1993). Major Rice producing countries of the world are China, India,

Bangladesh, Vietnam, Myanmar, Thailand, Japan, and Philippines, with China being the largest rice producing country (190 million tons) (Kole, 2011).

As population of the world increases, the demand for rice is also increasing enormously. The world population is supposed to be doubled by 2050 (van Beek *et al.*, 2010). The annual growth rate of world population is 1.3%, and 90% of this growth rate increase is taking place in developing countries of Africa, Latin America and Asia (Subudhi *et al.*, 2006). Cultivation can be increased via expanding the arable land. But the expansion of agricultural land seems to be impossible due to the rapid increase in industries, urbanization, and population. Cultivation area can be increased if the marginal soil is used; these lands have the potential to increase the food security (FOODS, 2008; Hill *et al.*, 2006; Robertson *et al.*, 2008). However, much of available agricultural soils and marginal lands are characterized by a number of abiotic stresses, like high acidity, uncontrolled flooding and the presence of toxic metals (e.g. iron and aluminum) that limit crop production in anaerobic and acid soils (Jia-en *et al.*, 2010).

Acidic soil is one of the main constraints in rice cultivation in many areas of the world (Bouton and Sumner, 1983). At low pH of the soil, aluminum (various species), Fe^{2+} and Mn^{2+} are solubilized into soil and that severely affect rice production (Kochian, 1995). Rice is cultivated by two methods, transplantation and direct seeded method. Direct seeded culture is a popular method as it gives higher yield and profit due to its low inputs and labour demand (Sushil Pandey, 2002). But in this method seeds of rice are directly exposed to Al^{3+} and Fe^{2+} toxicities in aerobic and acid soil that inhibit seed germination (Farooq *et al.*, 2011).

Fe^{2+} toxicity is a nutritional disorder which is caused by the excessive uptake of Fe^{2+} in low pH under an aerobic condition that damages different metabolic processes in rice plant (Nasr, 2013). The iron stress causes inhibition of seed germination, typical leaf-bronzing symptoms, stain edges and poorly developed dark brown root system. The reduction in height and dry matter has a correlation with rice production and ultimately affect the rice yield. Fe^{2+} toxicity has been reported in many Asian countries, including, Malaysia, Indonesia, Thailand, China, Philippines,

and India, where rice production is reduced by 12-100% depending on rice cultivars tolerance to Fe^{2+} and on the toxicity severity (Audebert and Sahrawat, 2008).

Aluminum (Al), when combined with acid soils, is very toxic to root system of crops that ultimately cause a reduction in vegetative growth and lower crop production (Kochian, 1995). It is reported that out of total 41,274,890 Km^2 of potentially cultivated land, 19,863,000 Km^2 (approximately half) is influenced by Al^{3+} toxicity (Bot *et al.*, 2000). In acidic soils, Al^{3+} toxicity reduces crop production 25-80% (Herrera-Estrella, 2003).

Rice is one of the model plants for molecular studies because its whole genome has been sequenced as it has a small genome size (430Mb). Ten known genome types (AA, BB, CC, BBCC, CCDD, EE, FF, GG, HHJJ, and HHKK) (Gramene, 2014) and 23 species of genus *Oryza* are identified. Fe^{2+} and Al^{3+} toxicity tolerance are complex traits in rice that influence the morphological and physiological factors. The advancement in molecular markers and high-density linkage maps has led to the identification of individual quantitative trait loci (QTLs) linked with, abiotic stress tolerance, insect resistance, disease resistance and several other agronomical traits in crop plants. Previous research has investigated the effects of Fe^{2+} and Al^{3+} toxicity on growth in rice (Ahsan *et al.*, 2007) and attempted to identify the tolerance mechanisms of rice to these metals, but genetic studies related to Fe^{2+} and Al^{3+} toxicity have not been documented well (Ahsan *et al.*, 2007; Odoemena, 1988). Breeding and genomic approaches to improve Fe^{2+} and Al^{3+} tolerance or resistance provides new opportunities to cultivate rice in acidic soil as well as in marginal lands to increase rice production. Therefore, it is necessary to study the genetic background of tolerance in rice to Fe^{2+} and Al^{3+} stress. Rice breeders are putting their efforts to develop the cultivators and hybrid with improved potential against Fe^{2+} and Al^{3+} toxicities (Sankar *et al.*, 2011). It would help plant breeders to develop new rice varieties capable of growing on marginal lands.

1.2 Problem Statement

Rapid increase in urbanization caused labour shortage that motivated farmers to shift from rice traditional transplantation cultivation to direct seeding cultivation as direct-seeding reduces labour demands and economic inputs. This method improves soil fertility, organic matter, soil filth and available moisture efficiency (Pandey and Velasco, 2002). But in problematic soil such as acidic soil, seeds are directly exposed to biotic and abiotic stresses as shown Figure 1.1. The paddy fields are reported to have high amounts/concentration of toxic elements like ferrous (Fe) aluminum (Al), copper (Cu), cobalt (Co), chromium (Cr), nickel (Ni), cadmium (Cd), and arsenic (As). Although some of these metals likewise Fe (20-1000mg/kg) and Al (0.2-30mg/g) are required by plant in traces. Their higher concentrations (e.g. Fe^{2+} (2000mg/kg and above and Al^{3+} above 30mg/g) can be toxic (Dufey *et al.*, 2014).




Figure 1.1: Problem statement diagram

Abiotic stresses are one of the main constraints in rice growth by direct seeded method in acidic soil. Among the abiotic stresses, Fe^{2+} and acid/ Al^{3+} severely inhibit rice seed germination as elaborated above in Figure 1.1. Germination of seeds are more sensitive than mature plants in response to Fe^{2+} and Al^{3+} stresses (Mehraban *et al.*, 2008). The main adverse effect of Fe^{2+} and Al^{3+} is the inhibition of root elongation that leads to inhibition of nutrient and water uptake, reduce leaf area, leaf bronzing, necrosis wilting and even plant death (Mehraban *et al.*, 2008).

Seed germination is influenced by Fe^{2+} and Al^{3+} in two ways (i) by their toxicities; and (ii) by their water uptake inhibition respectively (Karanner and Colville, 2011). Conventional strategies such as application of chemicals to alleviate soil pH are not appropriate strategies to reduce or avoid the toxicity of these metals. Alternatively, the effective way is to genetically improve rice tolerance through hybridization of (tolerant×sensitive) varieties with tolerance to Fe^{2+} and Al^{3+} stress (Jannink *et al.*, 2010).

Literature survey reveals that that genetically identification of QTLs (Quantitative Trait Loci) that linked with tolerance to Fe^{2+} and Al^{3+} toxicities for seed germination have not yet been carried out. Therefore, this research work was focused on the research gap to identify QTLs linked to Fe^{2+} and Al^{3+} toxicities tolerance in seed germination of *Oryza sativa*.

1.3 Research Objectives

- i) To develop a mapping population for genetic dissection of tolerance to Fe^{2+} and Al^{3+} toxicities at germination stage in *Indica* rice.
- ii) To optimize the tolerance limits of Fe^{2+} and Al^{3+} in parental lines for phenotyping of F_3 progenies.
- iii) To construct a molecular linkage map based on F_2 microsatellite marker data using MapMaker/EXP.
- iv) To identify the QTLs linked to seed germination tolerance against Fe^{2+} and Al^{3+} toxicities based on molecular linkage map and phenotypic data of F_3 progenies using WinQTL Cartographer.

1.4 Scope of Study

To determine the genetic basis of important agronomical traits, mapping populations construction is required. So in this study some selected rice genotypes were screened at germination stage under different doses of Fe^{2+} and Al^{3+} . After ANOVA and DMRT analysis, *Pokkali* has found as tolerant (Wu *et al.*, 2014) and

Pak Basmati as sensitive varieties (Javed *et al.*, 2011) among all. Mapping population has been constructed by crossing *Pak Basmati* (female parent) and *Pokkali* (male parent). The tolerance limits for Fe^{2+} and Al^{3+} toxicities were optimized where maximum significant differences have observed between the parental lines. F_3 progenies were evaluated for their tolerance to optimized levels of Fe^{2+} and Al^{3+} toxicities. The genotypic data of F_2 mapping population has used to construct a linkage map using microsatellites framework. WinQTL Cartographer version 2.5 software was used to identify the QTLs based on F_2 population, molecular linkage map and phenotypic traits of F_3 progenies.

1.5 Significance of Study

Since the demands to support and accelerate progress in breeding for novel traits has been increased, the plant breeding community needs to accurately analyze the increasingly large numbers of crop plants and their traits. In acidic soil, the solubility of Al^{3+} and Fe^{2+} increases that instantly increases translocation of these toxicant into rice crop, which result in germination inhibition and hinders plant establishment. The aim of this study is to provide quantitative analyses of plant genetically relevant for important traits that help plants better adapt to the acid soil and less agriculture input.

This research project has enhanced the genetic knowledge by identification of QTLs linked with tolerance to Fe^{2+} and Al^{3+} toxicities at the germination stage in *Indica* rice. Analysis and identification of putative QTL positions linked to QTLs tolerance to Fe^{2+} and Al^{3+} toxicities have explored the way in isolation and molecular characterization of QTLs via map based cloning. DNA markers that are tightly linked to agronomically main genes of tolerance to Fe^{2+} and Al^{3+} toxicities may be used as molecular tools for marker-assisted selection (MAS) in plant breeding. Subsequently, exploring the potential knowledge of QTLs linked to tolerance to Fe^{2+} and Al^{3+} toxicities detected on 12 linkage groups would help rice breeders to design breeding strategy to develop tolerant rice varieties to enhance the rice growth and yield.

REFERENCES

- Abdullah, Z., Khan, A. M., and Flowers, T. J. (2001). Causes of sterility in seed set of rice under salinity stress. *Journal of Agronomy and Crop Science*, 187(1), 25-32.
- Abdurakhmonov, I. Y., and Abdukarimov, A. (2008). Application of association mapping to understanding the genetic diversity of plant germplasm resources. *International Journal of Plant Genomics*, 2008, 18.
- Abedin, M. J., and Meharg, A. A. (2002). Relative toxicity of arsenite and arsenate on germination and early seedling growth of rice (*Oryza sativa* L.). *Plant and soil*, 243(1), 57-66.
- Adams, M. B., Burger, J. A., Jenkins, A. B., and Zelazny, L. (2000). Impact of harvesting and atmospheric pollution on nutrient depletion of eastern US hardwood forests. *Forest Ecology and Management*, 138(1-3), 301-319.
- Adrees, M., Ali, S., Rizwan, M., Zia-ur-Rehman, M., Ibrahim, M., Abbas, F., et al. (2015). Mechanisms of silicon-mediated alleviation of heavy metal toxicity in plants: a review. *Ecotoxicology and environmental safety*, 119, 186-197.
- Ahmad, A. R., and Nye, P. H. (1990). Coupled diffusion and oxidation of ferrous iron in soils. I. Kinetics of oxygenation of ferrous iron in soil suspension. *European Journal of Soil Science*, 41(3), 395-409.
- Ahmad, I., Akhtar, M. J., Ahmad, Z., and Jamil, A. (2012). Effect of cadmium on seed germination and seedling growth of four wheat (*Triticum aestivum* L.) cultivars. *Pakistan Journal of Botany*, 44(5), 1569-1574.
- Ahn, S.-N., Kim, D.-M., Lee, H.-S., Kang, J.-W., and Yun, Y.-T. (2016). *High-Density Mapping Reveals a Linkage of Quantitative Trait Loci for Grain-Weight and Spikelet Number in Rice*. Paper presented at the International Plant and Animal Genome Conference XXIV.

- Ahsan, N., Lee, D.-G., Lee, S.-H., Kang, K. Y., Lee, J. J., Kim, P. J., et al. (2007). Excess copper induced physiological and proteomic changes in germinating rice seeds. *Chemosphere*, 67(6), 1182-1193.
- Akinci, I. E., and Akinci, S. (2010). Effect of chromium toxicity on germination and early seedling growth in melon (*Cucumis melo* L.). *African Journal of Biotechnology*, 9(29), 4589-4594.
- Alamgir, A. N. M., and Akhter, S. (2009). Effects of aluminium (Al^{3+}) on seed germination and seedling growth of wheat (*Triticum aestivum* L.). *Bangladesh Journal of Botany*, 38(1), 1-6.
- Alamgir, M., and Hossain, M. K. (2005). Effect of pre-sowing treatments on germination and initial seedling development of Albizia saman in the nursery. *Journal of Forestry Research*, 16(3), 200-204.
- Allen, N. J., and Grisaffe, D. B. (2001). Employee commitment to the organization and customer reactions: Mapping the linkages. *Human Resource Management Review*, 11(3), 209-236.
- Almansouri, M., Kinet, J. M., and Lutts, S. (2001). Effect of salt and osmotic stresses on germination in durum wheat (*Triticum durum* Desf.). *Plant and Soil*, 231(2), 243-254.
- An, Y.-J. (2004). Soil ecotoxicity assessment using cadmium sensitive plants. *Environmental Pollution*, 127(1), 21-26.
- Ando, T., Yoshida, S., and Nishiyama, I. (1983). Nature of oxidizing power of rice roots. *Plant and Soil*, 72(1), 57-71.
- Aniol, A., and Gustafson, J. P. (1984). Chromosome location of genes controlling aluminum tolerance in wheat, rye, and triticale. *Canadian Journal of Genetics and Cytology*, 26(6), 701-705.
- Aniol, A., and Gustafson, J. P. (1989). Chapter 16: Genetics of tolerance in agronomic plants. In A. J. Shaw (Ed.), *Heavy Metal Tolerance in Plants: Evolutionary Aspects* (pp. 255-267). Google Books.
- Anjum, N. A., Sofo, A., Scopa, A., Roychoudhury, A., Gill, S. S., Iqbal, M., et al. (2015). Lipids and proteins—major targets of oxidative modifications in abiotic stressed plants. *Environmental Science and Pollution Research*, 22(6), 4099-4121.

- Ansari, M. U. R., Shaheen, T., Bukhari, S., and Husnain, T. (2015). Genetic improvement of rice for biotic and abiotic stress tolerance. *Turkish Journal of Botany*, 39(6), 911-919.
- Anuradha, K., Agarwal, S., Rao, Y. V., Rao, K. V., Viraktamath, B. C., and Sarla, N. (2012). Mapping QTLs and candidate genes for iron and zinc concentrations in unpolished rice of Madhukar×Swarna RILs. *Gene*, 508(2), 233-240.
- Armstrong, W., and Drew, M. C. (2002). Root Growth and Metabolism Under Oxygen Deficiency. In U. Kafkafi, Y. Waisel and A. Eshel (Eds.), *Plant Roots: The Hidden Half* (Vol. 3, pp. 729-761): CRC Press.
- Arnold, R. L. B., Fenner, M., and Edwards, P. J. (1991). Changes in germinability, ABA content and ABA embryonic sensitivity in developing seeds of *Sorghum bicolor* (L.) Moench. induced by water stress during grain filling. Netherlands: The Kluwer Academic Publishers.
- Arosio, P., and Levi, S. (2002). Ferritin, iron homeostasis, and oxidative damage^{and 1}
². *Free Radical Biology and Medicine*, 33(4), 457-463.
- Ashkani, S., Rafii, M., Rahim, H., and Latif, M. A. (2013). Genetic dissection of rice blast resistance by QTL mapping approach using an F3 population. *Molecular biology reports*, 40(3), 2503-2515.
- Audebert, A. (2006). Iron toxicity in rice—environmental conditions and symptoms. *Iron Toxicity in Rice-Based System in West Africa. WARDA, Cotonou*, 18-33.
- Audebert, A., and Fofana, M. (2008). Rice yield gap due to iron toxicity in West Africa. *Journal of Agronomy and Crop Science*, 195(1), 66-76.
- Audebert, A., and Sahrawat, K. L. (2008). Mechanisms for iron toxicity tolerance in lowland rice. *Journal of Plant Nutrition*, 208(38), 1877-1885.
- Ayaz, F. A., and Kadio lu, A. (1997). Effects of heavy metals (Zn, Cd, Cu, Hg) on the soluble protein bands of germinating *Lens Esculenta* L. seeds. *Turkish Journal of Botany*, 21(2), 85-88.
- Aydinalp, C., and Marinova, S. (2009). The effects of heavy metals on seed germination and plant growth on alfalfa plant (*Medicago sativa* L.). *Bulgarian Journal of Agricultural Science*, 15(4), 347-350.
- Bae, J., Benoit, D. L., and Watson, A. K. (2016). Effect of heavy metals on seed germination and seedling growth of common ragweed and roadside ground cover legumes. *Environmental Pollution*, 213, 112-118.

- Bakhat, H. F., Zia, Z., Fahad, S., Abbas, S., Hammad, H. M., Shahzad, A. N., et al. (2017). Arsenic uptake, accumulation and toxicity in rice plants: Possible remedies for its detoxification: A review. *Environmental Science and Pollution Research*, 1-17.
- Bassam, B. J., and Gresshoff, P. M. (2007). Silver staining DNA in polyacrylamide gels. *Nature protocols*, 2(11), 2649-2654.
- Basten, C. J., Weir, B. S., and Zeng, Z.-b. (1994). *Zmap—a QTL cartographer*. Paper presented at the 5th World Congress on Genetics Applied to Livestock Production: Computing Strategies and Software, Guelph, Ontario, Canada, 65-66.
- Basu, A., Basu, U., and Taylor, G. J. (1994). Induction of microsomal membrane proteins in roots of an aluminum-resistant cultivar of *Triticum aestivum* L. under conditions of aluminum stress. *Environmental and Plant Stress Physiology*, 104(3), 1007-1013.
- Becker, M., and Asch, F. (2005). Iron toxicity in rice — conditions and management concepts. *Journal of Plant Nutrition and Soil Science*, 168(4), 558-573.
- Bewley, J. D., Black, M., and Halmer, P. (Eds.). (2006) The encyclopedia of seeds (1 ed.). Wallingford United Kingdom: CAB International.
- Bhushan, L., Ladha, J. K., Gupta, R. K., Singh, S., Tirol-Padre, A., Saharawat, Y. S., et al. (2007). Saving of water and labor in a rice–wheat system with no-tillage and direct seeding technologies. *Agronomy Journal*, 99(5), 1288-1296.
- Bing, Z., Qi-Ming, D., Zhang, Q.-J., Jie-Qin, L., Shao-Ping, Y., Liang, Y.-S., et al. (2006). Analysis of segregation distortion of molecular markers in F₂ population of rice. *Acta Genetica Sinica*, 33(5), 449-457.
- Bitá, C., and Gerats, T. (2013). Plant tolerance to high temperature in a changing environment: scientific fundamentals and production of heat stress-tolerant crops. *Frontiers in plant science*, 4, 273.
- Bolan, N., Kunhikrishnan, A., Thangarajan, R., Kumpiene, J., Park, J., Makino, T., et al. (2014). Remediation of heavy metal (loid)s contaminated soils – To mobilize or to immobilize? *Journal of Hazardous Materials*, 266, 141-166.
- Bona, L., Carver, B. F., Wright, R. J., and Baligar, V. C. (1994). Aluminum tolerance of segregating wheat populations in acidic soil and nutrient solutions. *Communications in Soil Science & Plant Analysis*, 25(3-4), 327-339.

- Bot, A. J., Nachtergaele, F. O., and Young, A. (2000). *Land resource potential and constraints at regional and country levels* (Annual Report No. 9251044295). Rome: Food & Agriculture Organization (FAO). (F. a. A. O. F. o. t. U. Nations o. Document Number)
- Botstein, D., White, R. L., Skolnick, M., and Davis, R. W. (1980). Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *American journal of human genetics*, 32(3), 314-331.
- Bouton, J. H., and Sumner, M. E. (1983). Alfalfa, *Medicago sativa* L., in highly weathered, acid soils. *Plant and Soil: International Journal on Plant-Soil Relationships*, 74(3), 431-436.
- Brunner, I., and Sperisen, C. (2014). Aluminum exclusion and aluminum tolerance in woody plants. In B. Rewald, O. Falik, D. Godbold and S. Rachmilevitch (Eds.), *Ecophysiology of root systems-environment interaction* (Vol. 4, pp. 96-107): Frontiers in Plant Science.
- Budikova, S., and Durcekova, K. (2004). Aluminium accumulation in roots of Al-sensitive barley cultivar changes root cell structure and induces callose synthesis. *Biologia: Botany, Zoology and Cellular and Molecular Biology*, 59, 215-220.
- Cai, H., Zhou, Y., Xiao, J., Li, X., Zhang, Q., and Lian, X. (2009). Overexpressed glutamine synthetase gene modifies nitrogen metabolism and abiotic stress responses in rice. *Plant cell reports*, 28(3), 527-537.
- Camp, N. J. (2002). *Quantitative Trait Loci: Methods and Protocols* (Vol. 195). Totowa, New Jersey, United States: Humana Press Inc.
- Carson, M. A., and Morris, A. N. (2012). Germination of *Panicum virgatum* cultivars in a NaCl gradient. *BioOne Society (BIOS) Washington DC*, 83(3), 90-96.
- Chen, C., Zhou, P., Choi, Y. A., Huang, S., and Jr., F. G. G. (2006). Mining and characterizing microsatellites from citrus ESTs. *Theoretical and Applied Genetics*, 112(7), 1248-1257.
- Chen, Z., Zhu, Y. G., Liu, W. J., and Meharg, A. A. (2005). Direct evidence showing the effect of root surface iron plaque on arsenite and arsenate uptake into rice (*Oryza sativa* L.) roots. *New Phytologist*, 165(1), 91-97.
- Chitteti, B. R., and Peng, Z. (2007). Proteome and phosphoproteome differential expression under salinity stress in rice (*Oryza sativa*) roots. *Journal of Proteome Research*, 6(5), 1718-1727.

- Chmielewicz, K. M. (2002). Tutorial for QTX, software for genetic mapping of Mendelian markers and quantitative trait loci. In J. M. Meer, K. F. Manly and R. H. Cudmore (Eds.): Roswell Park Cancer Institute.
- Choudhary, A. K., Singh, D., and Kumar, J. (2011). A comparative study of screening methods for tolerance to aluminum toxicity in pigeonpea [*Cajanus cajan* (L.) Millspaugh]. *Australian Journal of Crop Science*, 5(11), 1419-1426.
- Chouhan, A., Sanjeeda, I., Maheshwari, R. S., and Bafna, A. (2012). Study of heavy metal accumulations and relative effects on total photosynthetic pigments in plant leaves growing in Pithampur industrial area sector -1, 2 and 3, India. *Journal of Environmental Research And Development*, 7(2A), 979-985.
- Chu, Z., Fu, B., Yang, H., Xu, C., Li, Z., Sanchez, A., et al. (2006). Targeting *xa13*, a recessive gene for bacterial blight resistance in rice. *Theoretical and Applied Genetics*, 112(3), 455-461.
- Clemens, S. (2006). Toxic metal accumulation, responses to exposure and mechanisms of tolerance in plants. *Biochimie*, 88(11), 1707-1719.
- Collard, B. C. Y., Jahufer, M. Z. Z., Brouwer, J. B., and Pang, E. C. K. (2005). An introduction to markers, quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: the basic concepts. *Euphytica: International Journal of Plant Breeding*, 142(1-2), 169-196.
- Cook, R. E. D. (1990). *PhD Thesis: Iron toxicity to wetland plants*. University of Sheffield, United Kingdom.
- Cranados, G., Pandey, S., and Ceballos, H. (1993). Response to Selection for Tolerance to Acid Soils in a Tropical Maize Population. *Alliance of Crop, Soil, and Environmental Science Societies (ACSESS)*, 33(5), 936-940.
- Cruz, R. P. d., and Milach, S. C. K. (2004). Cold tolerance at the germination stage of rice: methods of evaluation and characterization of genotypes. *Scientia Agricola: Journal of Science and Agriculture*, 61(1), 1-8.
- Datta, S. K. D. (1981). *Principles and Practices of Rice Production* (Vol. 1). New York: John Wiley & Sons Inc.
- Degenhardt, J., Larsen, P. B., Howell, S. H., and Kochian, L. V. (1998). Aluminum Resistance in the Arabidopsis Mutant *alr-104* Is Caused by an Aluminum-Induced Increase in Rhizosphere pH. *Whole Plant, Environmental and Stress Physiology*, 117(1), 19-27.

- Delhaize, E., Ma, J. F., and Ryan, P. R. (2012). Transcriptional regulation of aluminium tolerance genes. *Trends in plant science*, 17(6), 341-348.
- Deshmukh, R., Singh, A., Jain, N., Anand, S., Gacche, R., Singh, A., et al. (2010). Identification of candidate genes for grain number in rice (*Oryza sativa* L.). *Functional & integrative genomics*, 10(3), 339-347.
- Deshmukh, U. C., Saxena, R. R., Xalxo, M. S., Sharma, D., and Verulkar, S. B. (2013). Hybrid purity testing in rice (*Oryza sativa* L.) using microsatellite markers. *Electronic Journal of Plant Breeding*, 4(1), 1021-1026.
- Ding, D., Li, W., Song, G., Qi, H., Liu, J., and Tang, J. (2011). Identification of QTLs for arsenic accumulation in maize (*Zea mays* L.) using a RIL population. *PLoS One*, 6(10), e25646.
- Dixit, V., Pandey, V., and Shyam, R. (2001). Differential antioxidative responses to cadmium in roots and leaves of pea (*Pisum sativum* L. cv. *Azad*). *Journal of Experimental Botany*, 52(358), 1101-1109.
- Djedatin, G., Ndjiondjop, M.-N., Sanni, A., Lorieux, M., Verdier, V., and Ghesquiere, A. (2016). Identification of novel major and minor QTLs associated with *Xanthomonas oryzae* pv. *oryzae* (African strains) resistance in rice (*Oryza sativa* L.). *Rice*, 9(1), 1-10.
- Doerge, R. W. (2002). Mapping and analysis of quantitative trait loci in experimental populations. *Nature Reviews Genetics*, 3(1), 43-52.
- Dokku, P., Das, K. M., and Rao, G. J. N. (2013). Pyramiding of four resistance genes of bacterial blight in Tapaswini, an elite rice cultivar, through marker-assisted selection. *Euphytica: International Journal of Plant Breeding*, 192(1), 87-96.
- Doncheva, S., Amenós, M., Poschenrieder, C., and Barceló, J. (2005). Root cell patterning: a primary target for aluminium toxicity in maize. *Journal of experimental botany*, 56(414), 1213-1220.
- Dong, Y., Ogawa, T., Lin, D., Koh, H.-J., Kamiunten, H., Matsuo, M., et al. (2006). Molecular mapping of quantitative trait loci for zinc toxicity tolerance in rice seedling (*Oryza sativa* L.). *Field crops research*, 95(2), 420-425.
- Dufey, I., Draye, X., Lutts, S., Lorieux, M., Martinez, C., and Bertin, P. (2015a). Novel QTLs in an interspecific backcross *Oryza sativa* × *Oryza glaberrima* for resistance to iron toxicity in rice. *Euphytica*, 204(3), 609-625.

- Dufey, I., Hakizimana, P., Draye, X., Lutts, S., and Bertin, P. (2009). QTL mapping for biomass and physiological parameters linked to resistance mechanisms to ferrous iron toxicity in rice. *Euphytica*, 167(2), 143-160.
- Dufey, I., Mathieu, A.-S., Draye, X., Lutts, S., and Bertin, P. (2015b). Construction of an integrated map through comparative studies allows the identification of candidate regions for resistance to ferrous iron toxicity in rice. *Euphytica*, 203(1), 59-69.
- Dufey, I., S.Gheysens, A.Ingabire, Lutts, S., and P.Bertin. (2014). Silicon application in cultivated rices (*Oryza sativa* L and *Oryza glaberrima* Steud) alleviates iron toxicity symptoms through the reduction in iron concentration in the leaf tissue. *Journal of agronomy and crop science*, 200(2), 132-142.
- Edzesi, W. M., Dang, X., Liang, L., Liu, E., Zaid, I. U., and Hong, D. (2016). Genetic diversity and elite allele mining for grain traits in rice (*Oryza sativa* L.) by association mapping. *Frontiers in Plant Science Switzerland*, 7, 787.
- El-Hendawy, S. E., Hu, Y., Yakout, G. M., Awad, A. M., Hafiz, S. E., and Schmidhalter, U. (2005). Evaluating salt tolerance of wheat genotypes using multiple parameters. *European journal of agronomy*, 22(3), 243-253.
- El-Kherbawy, M., Angle, J. S., Heggo, A., and Chaney, R. L. (1989). Soil pH, rhizobia, and vesicular-arbuscular mycorrhizae inoculation effects on growth and heavy metal uptake of alfalfa (*Medicago sativa* L.). *Biology and Fertility of Soils*, 8(1), 61-65.
- Eswaran, H., Reich, P., and Beinroth, F. (1997). Global distribution of soils with acidity. *Plant-Soil interactions at low pH*, 1(1), 159-164.
- Fageria, N. K., Baligar, V. C., and Clark, R. B. (2006). *Physiology of crop production*. New York: Food Products Press.
- Fageria, V. D. (2001). Nutrient interactions in crop plants. *Journal of plant nutrition*, 24(8), 1269-1290.
- Fanping, M., and Guifang, L. (1998). The Effect of Acid Rain on Chemical Activities of Elements in Soil. *Journal Of Central South Forestry University China*, 1(1), 1-5.
- Faris, J. D., Laddomada, B., and Gill, B. S. (1998). Molecular Mapping of Segregation Distortion Loci in *Aegilops tauschii*. *Genetics Society of America*, 149(1), 319-327.

- Farooq, M., Basra, S. M. A., and Asad, S. A. (2008). Comparison of conventional puddling and dry tillage in rice - wheat system. *Paddy and Water Environment*, 6(4), 397-404.
- Farooq, M., Siddique, K. H. M., Rehman, H., Aziz, T., Lee, D.-J., and Wahid, A. (2011). Rice Direct Seeding: Experiences, Challenges and Opportunities. *Soil and Tillage Research*, 111(2), 87-98.
- Finatto, T. (2012). *Transcriptomic analysis of genes and LTR retrotransposons in rice (Oryza sativa L. ssp. japonica) in response to iron toxicity*. Unpublished Doctorate of Philosophy, Federal University of Pelotas, Brazil, Library Repository of Federal University of Pelotas, Brazil.
- FOODS, M. H. I. (2008). Food and agriculture organization of the United Nations (Publication., from Food and Agriculture Organization (FAO) of the United Nations, World Health Organization (WHO).
- Fran áková, H., Líšková, M., Boj ánská, T., and Mare ek, J. (2012). Germination Index as an Indicator of Malting Potential. *Czech Journal of Food Science*, 30(4).
- Frantzios, G., Galatis, B., and Apostolakis, P. (2001). Aluminium Effects on Microtubule Organization in Dividing Root-Tip Cells of *Triticum turgidum*. II. Cytokinetic Cells. *International Journal of Plant Research*, 114(2), 157-170.
- Frei, M., Tetteh, R. N., Razafindrazaka, A. L., Fuh, M. A., Wu, L.-B., and Becker, M. (2016). Responses of rice to chronic and acute iron toxicity: genotypic differences and biofortification aspects. *Plant and Soil*, 1-13.
- Fu, Z., Li, W., Xing, X., Xu, M., Liu, X., Li, H., et al. (2016). Genetic analysis of arsenic accumulation in maize using QTL mapping. *Scientific reports*, 6.
- Gangola, M. P., Khedikar, Y. P., Gaur, P. M., Båga, M., and Chibbar, R. N. (2013). Genotype and growing environment interaction shows a positive correlation between substrates of Raffinose Family Oligosaccharides (RFO) biosynthesis and their accumulation in chickpea (*Cicer arietinum* L.) seeds. *Journal of agricultural and food chemistry*, 61(20), 4943-4952.
- Gao, J.-S., Wu, N., Shen, Z.-L., Lv, K., Qian, S.-H., Guo, N., et al. (2016). Molecular cloning, expression analysis and subcellular localization of a Transparent Testa 12 ortholog in brown cotton (*Gossypium hirsutum* L.). *Gene*, 576(2), 763-769.

- Garvin, D. F., and Carver, B. F. (2003). Role of the genotype in tolerance to acidity and aluminum toxicity. *Handbook of Soil Acidity. Marcel Dekker, New York*, 387-406.
- Gramene. (2014). A comparative resource for plants Gramene. Retrieved 19 October 2016, from <http://gramene.org/>
- Gregorio, G. B., Senadhira, D., Mendoza, R. D., Manigbas, N. L., Roxas, J. P., and Guerta, C. Q. (2002). Progress in breeding for salinity tolerance and associated abiotic stresses in rice. *Journal of Field Crops Research*, 76(2), 91-101.
- Gross, J., Stein, R. J., Fett-Neto, A. G., and Fett, J. P. (2003). Iron homeostasis related genes in rice. *Journal of Genetics and Molecular Biology*, 26(4), 477-497.
- Gu, K., Sangha, J. S., Li, Y., and Yin, Z. (2008). High-resolution genetic mapping of bacterial blight resistance gene *Xa10*. *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 116(2), 155-163.
- Gu, K., Yang, B., Tian, D., Wu, L., Wang, D., Sreekala, C., et al. (2005). *R* gene expression induced by a type-III effector triggers disease resistance in rice. *Nature*, 435(7045), 1122-1125.
- Guo, L., Zhu, L., Xu, Y., Zeng, D., Wu, P., and Qian, Q. (2004). QTL analysis of seed dormancy in rice (*Oryza sativa* L.). *Euphytica: International Journal of Plant Breeding*, 140(3), 155-162.
- Gupta, N., Gaurav, S. S., and Kumar, A. (2013). Molecular basis of aluminium toxicity in plants: a review. *American Journal of Plant Sciences*, 2013.
- Gyawali, R., and Lekhak, H. D. (2006). Chromium tolerance of rice (*Oryza sativa* L.) cultivars from Kathmandu Valley, Nepal. *Scientific World Journal (Hindawi)*, 4(4), 102-108.
- Halwart, M., and Gupta, M. V. (2004). The Rice Field Ecosystem. In Matthias Halwart and M. V. Gupta (Eds.), *Culture of Fish in Rice Fields* (Vol. 1718, pp. 83). Rome, ItalyPenang, Malaysia: FAO:The WorldFish center.
- Harris, H. J. (1964). Delaware, United States Patent No. US 3150081 A. U. S. P. Office.
- Hashemi-Petroudi, S. H., Maibody, S. A. M. M., Nematzadeh, G. A., and Arzani, A. (2010). Semi-random PCR markers for DNA fingerprinting of rice hybrids

- and their corresponding parents. *African Journal of Biotechnology*, 9(7), 979-985.
- He, P., Li, J. Z., Zheng, X. W., Shen, L. S., Lu, C. F., Chen, Y., et al. (2001). Comparison of Molecular Linkage Maps and Agronomic Trait Loci between DH and RIL Populations Derived from the Same Rice Cross. *Crop Science: Alliance of Crop, Soil and Environmental Science Societies (ACSESS)*, 41(4), 1240-1246.
- Hede, A. R., Skovmand, B., and Cesati, J. L. (2001). *Acid soils and aluminum toxicity* (No. 9706480773): International Maize and Wheat Improvement Center (CIMMYT). (CIMMYT o. Document Number)
- Herrera-Estrella, L. (1999). Transgenic Plants for Tropical Regions: Some Considerations About Their Development and Their Transfer to the Small Farmer. *Proceedings of the National Academy of Sciences*, 96(11), 5978-5981.
- Herrera-Estrella, L. (2003, January. 30-31). *Case study: Use of biotechnology to increase food production on acid soils*. Paper presented at the Towards sustainable agriculture for developing countries: options from life sciences and biotechnologies, Brussels, Belgium, 5978-5981.
- Heun, M., Kennedy, A. E., Anderson, J. A., Lapitan, N. L. V., Sorrells, M. E., and Tanksley, S. D. (1991). Construction of a restriction fragment length polymorphism map for barley (*Hordeum vulgare*). *Genome: Canadian National Research Centre Research Press*, 34(3), 437-447.
- Hill, J., Nelson, E., Tilman, D., Polasky, S., and Tiffany, D. (2006). Environmental, economic, and energetic costs and benefits of biodiesel and ethanol biofuels. *Proceedings of the National Academy of sciences*, 103(30), 11206-11210.
- Holzschuh, M. J., Carlos, F. S., Carmona, F. d. C., Bohnen, H., and Anghinoni, I. (2014). Iron oxidation on the surface of adventitious roots and its relation to aerenchyma formation in rice genotypes. *Revista Brasileira de Ciência do Solo*, 38(1), 185-192.
- Hong, T. D., and Ellis, R. H. (1996). *A protocol to determine seed storage behaviour* (1 ed. Vol. 1). Department of Agriculture, The University of Reading, UK: International Plants genetic Resources Institute (IPGRI).

- Horst, W. J., Wang, Y., and Eticha, D. (2010). The role of the root apoplast in aluminium-induced inhibition of root elongation and in aluminium resistance of plants: A review. *Annals of Botany, Oxford Academic*, 106(1), 185-197.
- Huang, X. Q., Cloutier, S., Lycar, L., Radovanovic, N., Humphreys, D. G., Noll, J. S., et al. (2006). Molecular detection of QTLs for agronomic and quality traits in a doubled haploid population derived from two Canadian wheats (*Triticum aestivum* L.). *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 113(4), 753-766.
- Ikeda, N., Bautista, N. S., Yamada, T., Kamijima, O., and Ishii, T. (2001). Ultra-simple DNA extraction method for marker-assisted selection using microsatellite markers in rice. *Journal of Plant Molecular Biology Reporter*, 19(1), 27-32.
- Inglett, P. W., Reddy, K. R., and Corstanje, R. (2005). Anaerobic soils. *Encyclopedia of soils in the environment. Elsevier Science, Oxford, UK*, 1(1), 71-78.
- Inostroza-Blancheteau, C., Soto, B., Ulloa, P., Aquea, F., and Reyes-Díaz, M. (2008). Resistance mechanisms of aluminum (Al 3+) phytotoxicity in cereals: physiological, genetic and molecular bases. *Revista de la Ciencia del Suelo y Nutrición Vegetal*, 8(3), 57-71.
- IRRI. (1993). *Rice research in a time of change* (Annual Report): Los Banos, Laguna, the Philippines. (I. R. R. Institute o. Document Number)
- Iverson, C. E. (2014). Washington, DC Patent No. United States Patent Invision: U. S. P. a. T. Office.
- Jannink, J.-L., Lorenz, A. J., and Iwata, H. (2010). Genomic selection in plant breeding: from theory to practice. *Briefings in functional genomics*, 9(2), 166-177.
- Javed, M. A., Huyop, F. Z., Ishii, T., Samad, A. A., Mahmood, T., Haider, M. S., et al. (2013). Construction of microsatellite linkage map and detection of segregation distortion in indica rice (*Oryza sativa* L.). *Pakistan Journal of Botany*, 45, 2085-2092.
- Javed, M. A., Huyop, F. Z., Wagiran, A., and Salleh, F. M. (2011). Identification of QTLs for Morph-Physiological Traits Related to Salinity Tolerance at Seedling Stage in Indica Rice. *Procedia Environmental Sciences*, 8, 389-395.
- Jethani, I. (2000). Effect of Inhibitor on the Germination of Coriander and Cumin Seeds. *New Botanist: International journal of plant sciences*, 27(1/4), 25-28.

- Jia-en, W., Zhan-jun, K., Wei-feng, P., Rong-fu, R., Yan-hua, H., and Hong, L. (2010). Content Change and Forecast of Heavy Metal and pH Value in Soil for Jianshan Area, Northern Zhejiang Province from 1990 to 2008. *Geological Science and Technology Information China*, 1(1), 014.
- Johnson, A. A. T., Kyriacou, B., Callahan, D. L., Carruthers, L., Stangoulis, J., Lombi, E., et al. (2011). Constitutive Overexpression of the *OsNAS* Gene Family Reveals Single-Gene Strategies for Effective Iron- and Zinc-Biofortification of Rice Endosperm. *PLOS ONE*, 6(9), e24476.
- Joshi, B. H., and Modi, K. G. (2013). Screening and characterization of heavy metal resistant bacteria for its prospects in bioremediation of contaminated soil. *Journal of Environmental Research and Development Bhapol*, 7(4A), 1531-1538.
- Kamboj, R., Sharma, S., Vishwakarma, G. S., and Mittal, S. (2015). Arsenic induced physiological changes in improved varieties of rice grown in Malwa region of Punjab. *International Journal of Farm Sciences*, 5(4), 41-50.
- Kao, C.-H., Zeng, Z.-B., and Teasdale, R. D. (1999). Multiple Interval Mapping for Quantitative Trait Loci. *Genetics Journal of the Genetics Society of America*, 152(3), 1203-1216.
- Karanner, I., and Colville, L. (2011). Metals and seeds: Biochemical and molecular implications and their significance for seed germination. *Environmental and Experimental Botany*, 72(1), 93-105.
- Kearsey, M. J., and Pooni, H. S. (1998). *The Genetical Analysis of Quantitative Traits*: Stanley Thornes (Publishers) Ltd.
- Kendall, S., and Penfield, S. (2012). Maternal and zygotic temperature signalling in the control of seed dormancy and germination. *Seed Science Research*, 22(S1), S23-S29.
- Khan, M. R., and Khan, M. M. (2010). Effect of varying concentration of nickel and cobalt on the plant growth and yield of chickpea. *Australian Journal of Basic and Applied Sciences*, 4(1), 1036-1046.
- Khan, S. K., Iqbal, J., and Saeed, M. (2013). Comparative study of grain yield and biochemical traits of different rice varieties grown under saline and normal conditions. *Journal of Animal and Plant Sciences (JAPS)*, 23, 575-588.

- Knapp, A. K., and Anderson, J. E. (1980). Effect of heat on germination of seeds from serotinous lodgepole pine cones. *American Midland Naturalist Journal of University of Notre Dame, Indiana, United States*, 370-372.
- Kochian, L. V. (1995). Cellular mechanisms of aluminum toxicity and resistance in plants. *Annual review of plant biology*, 46(1), 237-260.
- Kochian, L. V., Pineros, M. A., and Hoekenga, O. A. (2005). The physiology, genetics and molecular biology of plant aluminum resistance and toxicity. In T. D. c. Hans Lambers (Ed.), *Root Physiology: From Gene to Function* (Vol. 4, pp. 175-195). Netherlands: Springer.
- Kole, C. (2011). *Wild Crop Relatives: Genomic and Breeding Resources: Cereals* (Vol. 1): Springer Science & Business Media, London New York.
- Konsten, C. J. M., Breemen, N. v., Suping, S., Aribawa, I. B., and Groenenberg, J. E. (1994). Effects of flooding on pH of rice-producing, acid sulfate soils in Indonesia. *Soil Science Society of America Journal*, 58(3), 871-883.
- Kothari, S. L., Davey, M. R., Lynch, P. T., Finch, R. P., and Cocking, E. C. (1993). Transgenic rice. *Transgenic plants Journal*, 2(1), 3-20.
- Koyama, M. L., Levesley, A., Koebner, R. M. D., Flowers, T. J., and Yeo, A. R. (2001). Quantitative Trait Loci for Component Physiological Traits Determining Salt Tolerance in Rice. *Journal of Plant Physiology: Environmental Stress and Adaptation*, 125(1), 406-422.
- Kreye, C., Bouman, B. A. M., Castaneda, A. R., Lampayan, R. M., Faronilo, J. E., Lactaoen, A. T., et al. (2009). Possible causes of yield failure in tropical aerobic rice. *Field crops research*, 111(3), 197-206.
- Lal, R. (2006). *Encyclopedia of soil science* (2006 ed. Vol. 2). United States of America: CRC Press.
- Lander, E. S., and Botstein, D. (1989). Mapping mendelian factors underlying quantitative traits using RFLP linkage maps. *Journal of Genetics: The Genetics Society of America*, 121(1), 185-199.
- Landsberg, E.-C. (1996). Hormonal regulation of iron-stress response in sunflower roots: a morphological and cytological investigation. *Protoplasma*, 194(1-2), 69-80.
- Lehmann, J., Rillig, M. C., Thies, J., Masiello, C. A., Hockaday, W. C., and Crowley, D. (2011). Biochar effects on soil biota—a review. *Soil Biology and Biochemistry*, 43(9), 1812-1836.

- Levesque, R. (2007). *SPSS Programming and Data Management: A Guide for SPSS and SAS Users*, (2007), SPSS Inc (4 ed.). Chicago Illinois: SPSS Inc.
- Li, W., Khan, M. A., Yamaguchi, S., and Kamiya, Y. (2005). Effects of heavy metals on seed germination and early seedling growth of *Arabidopsis thaliana*. *International Journal of Plant Growth Regulation*, 46(1), 45-50.
- Li, W., Tang, D., Wu, W., Lu, H., and Worland, A. J. (1999). A molecular map based on an *indica/indica* recombinant inbred population and its comparison with an existing map derived from *indica/japonica* cross in rice. *Chinese Journal of Rice Science (Zhongguo shuidao kexue)*, 14(2), 71-78.
- Li, X., Ma, H., Jia, P., Wang, J., Jia, L., Zhang, T., et al. (2012). Ecotoxicology and Environmental Safety Responses of seedling growth and antioxidant activity to excess iron and copper in *Triticum aestivum* L. *Ecotoxicology and Environmental Safety*, 86, 47-53.
- Lima, M., deMiranda, J. B., and Furlani, P. R. (1995). Diallel cross among inbred lines of maize differing in aluminum tolerance. *Brazilian Journal of Genetics*, 18(4), 579-584.
- Lin, Y.-F., and Aarts, M. G. M. (2012). The molecular mechanism of zinc and cadmium stress response in plants. *Cellular and molecular life sciences*, 69(19), 3187-3206.
- Lincoln, S. E., Daly, M. J., and Lander, E. S. (1993). *Mapping genes controlling quantitative traits with MAPMAKER/QTL 1.1* (Research Technical Report). Cambridge, USA: Whitehead Institute for Biomedical. (C. C. Whitehead Institute o. Document Number)
- Lincoln, S. E., and Newburg, L. (1987). MAPMAKER: an interactive computer program for constructing genetic linkage maps of experimental and natural populations. *Plant Molecular Biology Report*, 7(1), 116-128.
- Liphadzi, M. S., and Kirkham, M. B. (2006). Physiological Effects of Heavy Metals on Plant Growth and Function. In B. Huang (Ed.), *Plant-environment interactions* (3 ed., pp. 243-270). United States of America (USA): CRC Press, Taylor & Francis.
- Liu, B. H. (1997). QTL Mapping:Single-Marker analysis. In B. H. Liu (Ed.), *Statistical genomics: linkage, mapping, and QTL analysis* (2 ed., pp. 387-416). United states of America (USA): CRC press.

- Liu, G., Bernhardt, J. L., Jia, M. H., Wamishe, Y. A., and Jia, Y. (2008). Molecular characterization of the recombinant inbred line population derived from a *japonica-indica* rice cross. *Euphytica*, 159(1-2), 73-82.
- Liu, W. J., Zhu, Y. G., Smith, F. A., and Smith, S. E. (2004). Do iron plaque and genotypes affect arsenate uptake and translocation by rice seedlings (*Oryza sativa* L.) grown in solution culture? *Journal of Experimental Botany*, 55(403), 1707-1713.
- Liu, X., Zhang, S., Shan, X., and Zhu, Y.-G. (2005). Toxicity of arsenate and arsenite on germination, seedling growth and amylolytic activity of wheat. *Chemosphere: International Multidisciplinary Review Journal* 61(2), 293-301.
- Llugany, M., Poschenrieder, C., and Barceló, J. (1995). Monitoring of aluminium-induced inhibition of root elongation in four maize cultivars differing in tolerance to aluminium and proton toxicity. *Physiologia Plantarum: international journal of plant physiology*, 93(2), 265-271.
- Lorieux, M., Perrier, X., Goffinet, B., Lanaud, C., and León, D. G. D. (1995). Maximum-likelihood models for mapping genetic markers showing segregation distortion. 2. F₂ populations. *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 90(1), 81-89.
- Lucena, J. J. (2000). Effects of bicarbonate, nitrate and other environmental factors on iron deficiency chlorosis: A review. *Journal of Plant Nutrition*, 23(11-12), 1591-1606.
- Lui, D., and Wang, Y. (2002). *Arsenic. In Diagnostic Criteria for Plants and Soils*. Berkeley, California: University of California press. Document Number)
- Lyttle, T. W. (1991). Segregation distorters. *Annual review of genetics*, 25(1), 511-581.
- M.Sudharani, S.Rao, P., and Rao, L. V. S. (2014). Identification of SSR Markers for Testing of Hybridity and Seed Genetic Purity in Maize (*Zea Mays* L.) *International Journal of Science and Research (IJSR)* (Vol. 3, pp. 92-95). Rajendranagar, Hyderabad, India: Seed Research and Technology Centre.
- Ma, J. F., Shen, R., Zhao, Z., Wissuwa, M., Takeuchi, Y., Ebitani, T., et al. (2002). Response of rice to Al stress and identification of quantitative trait loci for Al tolerance. *International Journal of Plant and Cell Physiology*, 43(6), 652-659.

- Ma, J. F., Taketa, S., and Yang, Z. M. (2000). Aluminum Tolerance Genes on the Short Arm of Chromosome 3R Are Linked to Organic Acid Release in Triticale. *Journal of Plant Physiology: Environmental and Stress Physiology*, 122(3), 687-694.
- Ma, Z., and Miyasaka, S. C. (1998). Oxalate Exudation by Taro in Response to Al. *Journal of Plant Physiology: Environmental and Stress Physiology*, 118(3), 861-865.
- Maccaferri, M., Sanguineti, M. C., Corneti, S., Ortega, J. L. A., Salem, M. B., Bort, J., et al. (2008). Quantitative trait loci for grain yield and adaptation of durum wheat (*Triticum durum* Desf.) across a wide range of water availability. *Genetics*, 178(1), 489-511.
- Maheaswari, R., Kshirsagar, J. T., and Lavanya, N. (2016). Polymerase chain reaction: A molecular diagnostic tool in periodontology. *Journal of Indian Society of Periodontology*, 20(2), 128.
- Manly, K. F., Jr., R. H. C., and Meer, J. M. (2001). Map Manager QTX, cross-platform software for genetic mapping. *Mammalian Genome Journal of International Mammalian Genome Society*, 12(12), 930-932.
- Mapping, M., Aluminum, O. F., Nguyen, B. A. Y. D. U. Y., Submitted, D., Fulfillment, P., Of, D., et al. (2001). MOLECULAR MAPPING OF ALUMINUM.
- McCouch, S. R., Chen, X., Panaud, O., Temnykh, S., Xu, Y., Cho, Y. G., et al. (1997). Microsatellite Marker Development, Mapping and Applications In Rice Genetics and Breeding. *International Journal of Plant Molecular Biology*, 35(1-2), 89-99.
- McCouch, S. R., Temnykh, S., Lukashova, A., Coburn, J., DeClerck, G., Cartinhour, S., et al. (2001). Microsatellite Markers in Rice: Abundance, Diversity, and Applications. In G. S. Khush, D. S. Brar and B. Hardy (Eds.), *Rice Genetics IV* (Vol. 1, pp. 117-135): International Rice Research Institute (IRRI) Manila, Philippines.
- Meharg, A., and Marschner, H. P. (2012). *Marschner's Mineral Nutrition of Higher Plants* (Vol. 48). Wallingford, UK: Cambridge University Press.
- Meharg, A. A., and Rahman, M. M. (2003). Arsenic Contamination of Bangladesh Paddy Field Soils: Implications for Rice Contribution to Arsenic

- Consumption. *International Journal of Environmental Science and Technology*, 37(2), 229-234.
- Mehraban, P., Zadeh, A. A., and Sadeghipour, H. R. (2008). Iron toxicity in rice (*Oryza sativa* L.), under different potassium nutrition. *Asian Journal of Plant Sciences*, 7(3), 251-259.
- Miftahudin, A., Chikmawati, T., Ross, K., Scoles, G. J., and Gustafson, J. P. (2005). Targeting the aluminum tolerance gene *Alt3* region in rye, using rice/rye micro-colinearity. *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 110(5), 906-913.
- Miles, C., and Wayne, M. (2008). Quantitative trait locus (QTL) analysis. *Nature Education and Science*, 1(1), 208.
- Miura, K., Ashikari, M., and Matsuoka, M. (2011). The role of QTLs in the breeding of high-yielding rice. *Trends in plant science*, 16, 319-326.
- Miyasaka, S. C., Buta, J. G., Howell, R. K., and Foy, C. D. (1991). Mechanism of aluminum tolerance in snapbeans root exudation of citric acid. *Journal of Plant Physiology: Environmental and Stress Physiology*, 96(3), 737-743.
- Moncada, P., Martinez, C. P., Borrero, J., Châtel, M., Jr., H. G., Guimaraes, E., et al. (2001). Quantitative trait loci for yield and yield components in an *Oryza sativa*×*Oryza rufipogon* BC₂F₂ population evaluated in an upland environment. *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 102(1), 41-52.
- Morita, A., Yanagisawa, O., Takatsu, S., Maeda, S., and Hiradate, S. (2008). Mechanism for the detoxification of aluminum in roots of tea plant (*Camellia sinensis* (L.) Kuntze). *Phytochemistry International Journal*, 69(1), 147-153.
- Muhrizal, S., Shamshuddin, J., Fauziah, I., and Husni, M. A. H. (2006). Changes in iron-poor acid sulfate soil upon submergence. *Geoderma: The Global Journal of Soil Science*, 131(1), 110-122.
- Mullis, K., Faloona, F., Scharf, S., Saiki, R. I., Horn, G., and Erlich, H. (1986). *Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction*. Paper presented at the Cold Spring Harbor symposia on quantitative biology, 263-273.
- Munongo, S., and Shallone, C. K. (2013). Estimating the role of agricultural technologies in improving rural household welfare: A case of Masvingo. *Russian Journal of Agricultural and Socio-Economic Sciences*, 14(2).

- Munzuroglu, O., and Geckil, H. (2002). Effects of metals on seed germination, root elongation, and coleoptile and hypocotyl growth in *Triticum aestivum* and *Cucumis sativus*. *Archives of Environmental Contamination and Toxicology*, 43(2), 203-213.
- Murthy, U. M. N., Kumar, P. P., and Sun, W. Q. (2003). Mechanisms of seed ageing under different storage conditions for *Vigna radiata* (L.) Wilczek: lipid peroxidation, sugar hydrolysis, Maillard reactions and their relationship to glass state transition. *Journal of experimental Botany*, 54(384), 1057-1067.
- Nagata, T., Hayatsu, M., and Kosuge, N. (1992). Identification of aluminium forms in tea leaves by ²⁷Al NMR. *International Journal of Phytochemistry*, 31(4), 1215-1218.
- Nasr, N. (2013). Germination and seedling growth of maize (*Zea mays* L.) seeds in toxicity of aluminum and nickel. *Merit Research Journal of Environmental Science and Toxicology*, 1(5), 110-113.
- Nayak, A. K., Raja, R., Rao, K. S., Shukla, A. K., Mohanty, S., Shahid, M., et al. (2015). Effect of fly ash application on soil microbial response and heavy metal accumulation in soil and rice plant. *Ecotoxicology and environmental safety*, 114, 257-262.
- Nelson, J. C. (1997). QGENE: software for marker-based genomic analysis and breeding. *Molecular breeding*, 3(3), 239-245.
- Newtona, L. A., Chiltona, N. B., Beveridgea, I., Hosteb, H., Nansenc, P., and Gasser, R. B. (1998). Genetic markers for stronglyid nematodes of livestock defined by PCR-based restriction analysis of spacer rDNA. *Acta tropica: international journal on infectious diseases*, 69(1), 1-15.
- Nguyen, B. D., Brar, D. S., Bui, B. C., Nguyen, T. V., Pham, L. N., and Nguyen, H. T. (2003). Identification and mapping of the QTL for aluminum tolerance introgressed from the new source, ORYZA RUFIPOGON Griff., into indica rice (*Oryza sativa* L.). *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 106(4), 583-593.
- Nguyen, V., Nguyen, B., Sarkarung, S., Martinez, C., Paterson, A., and Nguyen, H. (2002). Mapping of genes controlling aluminum tolerance in rice: Comparison of different genetic backgrounds. *Molecular Genetics and Genomics*, 267(6), 772-780.

- Nguyen, V. T., Burow, M. D., Nguyen, H. T., Le, B. T., Le, T. D., and Paterson, A. H. (2001). Molecular mapping of genes conferring aluminum tolerance in rice (*Oryza sativa* L.). *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 102(6-7), 1002-1010.
- Odoemena, C. S. (1988). Breaking of seed coat dormancy in a medicinal plant *Tetrapleura tetraptera* (Schum & Thonn). *The Journal of Agricultural Science*, 111(02), 393-394.
- Oka, H.-I. (1988). *Origin of Cultivated Rice* (1st Edition ed. Vol. 14). Tokyo Japan: Japan Science Societies Press, Elsevier.
- Ologundudu, A. F., Adelusi, A. A., and Adekoya, K. P. (2013). Effect of Light Stress on Germination and Growth Parameters of *Corchorus olitorius*, *Celosia argentea*, *Amaranthus cruentus*, *Abelmoschus esculentus* and *Delonix regia*. *Notulae Scientia Biologicae Journal*, 5(4), 468.
- Ooijen, J. W. V. (1999). LOD significance thresholds for QTL analysis in experimental populations of diploid species. *Journal of Heredity: Oxford Academic*, 83(5), 613-624.
- Pallavi, H. M., Gowda, R., Shadakshari, Y. G., Bhanuprakash, K., and Vishwanath, K. (2011). Identification of SSR markers for hybridity and seed genetic purity testing in sunflower (*Helianthus annuus* L.). *HELIA: International Scientific Journal of the F.A.O Eeuropean Cooperative Research Network on Sunflower and the International Sunflower Association*, 34(54), 59-66.
- Pandey, N., and Sharma, C. P. (2002). Effect of heavy metals Co^{2+} , Ni^{2+} and Cd^{2+} on growth and metabolism of cabbage. *International Journal of Plant Science*, 163(4), 753-758.
- Pandey, S., and Velasco, L. (2002). Economics of direct seeding in Asia: Patterns of adoption and research priorities. In S. Pandey, M. Mortimer, L. Wade, T. P. Tuong, K. Lopez and B. Hardy (Eds.), *Direct seeding: research strategies and opportunities*. (Vol. 1, pp. 3-14): International Rice Research Institute, Los Baños (Philippines).
- Pandey, S., and Velasco, L. E. (1999). Economics of alternative rice establishment methods in Asia: a strategic analysis. *Social Sciences Division Discussion Paper, International Rice Research Institute, Los Banos, Philippines*, 1(1), 12-18.

- Passos, L. P., Köpp, M. M., and Léo, F. J. S. (2012). Performance of tetraploid alfalfa genotypes as exposed to aluminum toxicity. *Journal of Agricultural Sciences*, 3(2), 1-11.
- Paterson, A. H. (1996). Making genetic maps. *Genome mapping in plants*, 194.
- Paterson, A. H., Damon, S., Hewitt, J. D., Zamir, D., Rabinowitch, H. D., Lincoln, S. E., et al. (1991a). Mendelian factors underlying quantitative traits in tomato: comparison across species, generations, and environments. *Journal of Genetics: The Genetics Society of America*, 127(1), 181-197.
- Paterson, A. H., Tanksley, S. D., and Sorrells, M. E. (1991b). DNA markers in plant improvement. *Journal of Advances in agronomy*, 46(1), 39-90.
- Pellet, D. M., Grunes, D. L., and Kochian, L. V. (1995). Organic acid exudation as an aluminum-tolerance mechanism in maize (*Zea mays* L.). *Planta: International Journal of Plant Biology*, 196(4), 788-795.
- Peralta, J. R., Gardea-Torresdey, J. L., Tiemann, K. J., Gomez, E., Arteaga, S., Rascon, E., et al. (2001). Uptake and effects of five heavy metals on seed germination and plant growth in alfalfa (*Medicago sativa* L.). *Bulletin of Environmental Contamination and Toxicology*, 66(6), 727-734.
- Poschenrieder, C., Tolra, R., Hajiboland, R., Arroyave, C., and Barceló, J. (2015). Mechanisms of Hyper-resistance and Hyper-tolerance to Aluminum in Plants. In *Aluminum Stress Adaptation in Plants* (pp. 81-98): Springer.
- Powell, W., Morgante, M., Andre, C., Hanafey, M., Vogel, J., Tingey, S., et al. (1996). The comparison of RFLP, RAPD, AFLP and SSR (microsatellite) markers for germplasm analysis. *Molecular breeding Journal: New Strategies in Plant Improvement*, 2(3), 225-238.
- Pradhan, S. K., Nayak, D. K., Mohanty, S., Behera, L., Barik, S. R., Pandit, E., et al. (2015). Pyramiding of three bacterial blight resistance genes for broad-spectrum resistance in deepwater rice variety, Jalmagna. *Rice*, 8(1), 1-14.
- Purcell, L. C., Keisling, T. C., and Sneller, C. H. (2002). Soybean yield and water extraction in response to deep tillage and high soil aluminum. *Communications in soil science and plant analysis*, 33(19-20), 3723-3735.
- Qi, X., Pittaway, T. S., Lindup, S., Liu, H., Waterman, E., Padi, F. K., et al. (2004). An integrated genetic map and a new set of simple sequence repeat markers for pearl millet, *Pennisetum glaucum*. *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 109(7), 1485-1493.

- Qiu, F., Zheng, Y., Zhang, Z., and Xu, S. (2007). Mapping of QTL Associated with Waterlogging Tolerance during the Seedling Stage in Maize. *Annals of Botany: Oxford Academic*, 99(6), 1067-1081.
- Rajpurohit, D., Kumar, R., Kumar, M., Paul, P., Awasthi, A., Basha, P. O., et al. (2011). Pyramiding of two bacterial blight resistance and a semidwarfing gene in Type 3 Basmati using marker-assisted selection. *Euphytica: International Journal of Plant Breeding*, 178(1), 111-126.
- Ranal, M. A., and Santana, D. G. d. (2006). How and why to measure the germination process? *Brazilian Journal of Botany*, 29(1), 1-11.
- Ranawake, A. L., Manangkil, O. E., Yoshida, S., Ishii, T., Mori, N., and Nakamura, C. (2014). Mapping QTLs for cold tolerance at germination and the early seedling stage in rice (*Oryza sativa* L.). *Biotechnology & Biotechnological Equipment*, 28(6), 989-998.
- Rangel, A. F., Rao, I. M., and Horst, W. J. (2009). Intracellular distribution and binding state of aluminum in root apices of two common bean (*Phaseolus vulgaris*) genotypes in relation to Al toxicity. *Physiologia Plantarum*, 135(2), 162-173.
- Rao, I. M., Zeigler, R. S., Vera, R., and Sarkarung, S. (1993). Selection and breeding for acid-soil tolerance in crops. *Journal of BioScience*, 43(7), 454-465.
- Rees, F., Germain, C., Sterckeman, T., and Morel, J.-L. (2015). Plant growth and metal uptake by a non-hyperaccumulating species (*Lolium perenne*) and a Cd-Zn hyperaccumulator (*Noccaea caerulea*) in contaminated soils amended with biochar. *Plant and soil*, 395(1-2), 57-73.
- Robertson, G. P., Dale, V. H., Doering, O. C., Hamburg, S. P., Milillo, J. M., Wander, M. M., et al. (2008). Agriculture - Sustainable biofuels Redux. *United States Journal of Science*, 322(5898).
- Rodríguez, M., Canales, E., and Borrás-Hidalgo, O. (2005). Molecular aspects of abiotic stress in plants. *Biotecnología Aplicada*, 22(1), 1-10.
- Rout, G., Samantaray, S., and Das, P. (2001). Aluminium Toxicity in Plants: A Review. *Agronomie: International Journal in Agriculture and Environment*, 21(1), 3-21.
- Rout, G. R., and Sahoo, S. (2015). Role of iron in plant growth and metabolism. *Reviews in Agricultural Science*, 3(1), 1-24.

- Ryan, P. R., and Delhaize, E. (2010). The convergent evolution of aluminium resistance in plants exploits a convenient currency. *Functional Plant Biology*, 37(4), 275-284.
- Ryan, P. R., Delhaize, E., and Jones, D. L. (2001). Function and mechanism of organic anion exudation from plant roots. *Annual review of plant biology*, 52(1), 527-560.
- Ryan, P. R., Ditomaso, J. M., and Kochian, L. V. (1993). Aluminium toxicity in roots: an investigation of spatial sensitivity and the role of the root cap. *Journal of Experimental Botany*, 44(2), 437-446.
- Sahrawat, K. L. (2005). Iron toxicity in wetland rice and the role of other nutrients. *Journal of Plant Nutrition*, 27(8), 1471-1504.
- Sankar, P. D., Saleh, M. A. A. M., and Selvaraj, C. I. S. (2011). Rice breeding for salt tolerance. *Research in Biotechnology*, 2(2), 1-10.
- Scott, S. J., Jones, R. A., and Williams, W. A. I. (1984). Review of data analysis methods for seed germination. *International Journal of Crop sciences*, 24(6), 1192-1199.
- Selvakumar, P., Ravikesavan, R., Gopikrishnan, A., Thiyagu, K., Preetha, S., and Boopathi, N. M. (2010). Genetic purity analysis of cotton (*Gossypium spp.*) hybrids using SSR markers. *Seed Science and Technology*, 38(2), 358-366.
- Semagn, K., Bjørnstad, Å., and Ndjiondjop, M. N. (2006). Principles, requirements and prospects of genetic mapping in plants. *African Journal of Biotechnology*, 5(25), 2569-2587.
- Septiningsih, E. M., Prasetyono, J., Lubis, E., Tai, T. H., Tjubaryat, T., Moeljopawiro, S., et al. (2003). Identification of quantitative trait loci for yield and yield components in an advanced backcross population derived from the *Oryza sativa* L. variety IR64 and the wild relative *O. rufipogon*. *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 107(8), 1419-1432.
- Seregin, I. V., and Kozhevnikova, A. D. (2005). Distribution of cadmium, lead, nickel, and strontium in imbibing maize caryopses. *Russian Journal of Plant Physiology*, 52(4), 565-569.
- Shah, S. S., Mohammad, F., Shafi, M., Bakht, J., and Zhou, W. (2011). Effects of cadmium and salinity on growth and photosynthesis parameters of *Brassica* species. *Pakistan Journal of Botany*, 43(1), 333-340.

- Shamshuddin, J., Azura, A. E., Shazana, M. A. R. S., Fauziah, C. I., Panhwar, Q. A., and Naher, U. A. (2014). Properties and Management of Acid Sulphate Soils in Southeast Asia for Sustainable Cultivation of Rice, Oil Palm, and Cocoa. In D. L. Sparks (Ed.), *Advances in Agronomy* (Vol. 124, pp. 91-142). United States of Americas: Academic Press Publication, Elsevier.
- Shamshuddin, J., Elisa, A. A., Shazana, M. A. R. S., and Fauziah, I. C. (2013). Rice defense mechanisms against the presence of excess amount of Al^{3+} and Fe^{2+} in the water. *Australian Journal of Crop Science*, 7(3), 314.
- Sharma, N., and Kaur, R. (2014). MAPMAKER/EXP Version 3.0b: Though Used Since Time Immemorial Still Difficult for Beginners to Start with Map Construction for the Genetic Studies. *Journal of Translational Proteomics Research*, 1(1), 17-27.
- Shimizu, A. (2009). QTL analysis of genetic tolerance to iron toxicity in rice (*Oryza sativa* L.) by quantification of bronzing score. *Journal of New Seeds*, 10(3), 171-179.
- Shimizu, A., Guerta, C. Q., Gregorio, G. B., Kawasaki, S., and Ikehishi, H. (2005). QTLs for nutritional contents of rice seedlings (*Oryza sativa* L.) in solution cultures and its implication to tolerance to iron-toxicity. *Plant and Soil: International Journal on Plant-Soil Relationships*, 275(1-2), 57-66.
- Shinohara, M., Aoyama, C., Fujiwara, K., Watanabe, A., Ohmori, H., Uehara, Y., et al. (2011). Microbial mineralization of organic nitrogen into nitrate to allow the use of organic fertilizer in hydroponics. *Soil science and plant nutrition*, 57(2), 190-203.
- Sibov, S. T., Gaspar, M., Silva, M. J., Ottoboni, L. M. M., Arruda, P., and Souza, A. P. (1999). Two genes control aluminum tolerance in maize: Genetic and molecular mapping analyses. *Genome: Canadian National Research Centre Research Press*, 42(3), 475-482.
- Silva, S. (2012). Aluminium toxicity targets in plants. *Journal of Botany, Hindawi*, 2012, 1-8.
- Singh, Y., Singh, G., Johnson, D., and Mortimer, M. (2005). Changing from transplanted rice to direct seeding in the rice-wheat cropping system in India. In K. Toriyama, K. L. Heong and B. Hardy (Eds.), *Rice Is Life: Scientific Perspectives for the 21st Century* (Vol. 1, pp. 198-201): Proceedings of the World Rice Research Conference, Tsukuba, Japan.

- Siqueira-Silva, A. I., Silva, L. C. d., Azevedo, A. A., and Oliva, M. A. (2012). Iron plaque formation and morphoanatomy of roots from species of restinga subjected to excess iron. *Ecotoxicology and environmental safety*, 78(1), 265-275.
- Snowden, R. E. D., and Wheeler, B. D. (1993). Iron toxicity to fen plant species. *Journal of Ecology*, 1(1), 35-46.
- Soller, M., Brody, T., and Genizi, A. (1979). The expected distribution of marker-linked quantitative effects in crosses between inbred lines. *Journal of Heredity: Oxford Academic*, 43(1), 179-190.
- Song, X.-J., and Ashikari, M. (2008). Toward an optimum return from crop plants. *Rice: Applied Plant Science Journal*, 1(2), 135-143.
- Sreenivasulu, N., Sopory, S., and Kishor, P. K. (2007). Deciphering the regulatory mechanisms of abiotic stress tolerance in plants by genomic approaches. *Gene*, 388(1), 1-13.
- Srivastava, J., Nautiyal, A. R., Chandra, H., and Kalra, S. J. S. (2012). Response of C₃ and C₄ Plant Systems Exposed to Heavy Metals for Phytoextraction at Elevated Atmospheric CO₂ and at Elevated Temperature. In D. J. Srivastava (Ed.), *Environmental Contamination* (pp. 1-16). Shanghai China: INTECH Publisher Inc.
- Stein, R. J., Duarte, G. L., Spohr, M. G., Lopes, S. I. G., and Fett, J. P. (2009). Distinct physiological responses of two rice cultivars subjected to iron toxicity under field conditions. *Annals of Applied Biology*, 154(2), 269-277.
- Stockwell, C. A., Hendry, A. P., and Kinnison, M. T. (2003). Contemporary evolution meets conservation biology. *Trends in Ecology & Evolution*, 18(2), 94-101.
- Subudhi, P. K., Sasaki, T., and Khush, G. S. (2006). 1 Rice. In T. Sasaki (Ed.), *Rice* (Vol. 2, pp. 1-9). Verlag Berlin Heidelberg: Springer.
- Sundaram, R. M., Vishnupriya, M. R., Biradar, S. K., Laha, G. S., Reddy, G. A., Rani, N. S., et al. (2008). Marker assisted introgression of bacterial blight resistance in *Samba Mahsuri*, an elite *indica* rice variety. *Euphytica: International Journal of Plant Breeding*, 160(3), 411-422.
- Susanto, U., Aswidinnoor, H., Koswara, J., Setiawan, A., Lopena, V., Torizo, L., et al. (2008). QTL mapping of yield, yield components, and morphological

- traits in rice (*Oryza sativa* L.) using SSR marker. *Jurnal Agronomi Indonesia (Indonesian Journal of Agronomy)*, 36(3).
- Sushil Pandey. (2002). *Direct seeding: research strategies and opportunities* (Vol. 1). Metro Manila, Phillippines: International Rice Research Institute (IRRI).
- Tabuchi, A., and Matsumoto, H. (2001). Changes in cell-wall properties of wheat (*Triticum aestivum*) roots during aluminum-induced growth inhibition. *Physiologia Plantarum Journal*, 112(3), 353-358.
- Takahashi, R., Ishimaru, Y., Senoura, T., Shimo, H., Ishikawa, S., Arao, T., et al. (2011). The OsNRAMP₁ iron transporter is involved in Cd accumulation in rice. *Journal of Experimental Botany*, 62(14), 4843-4850.
- Tautz, D. (1989). Hypervariability of simple sequences as a general source for polymorphic DNA markers. *Journal of Nucleic acids research*, 17(16), 6463-6471.
- Temnykh, S., Park, W. D., Ayres, N., Cartinhour, S., Hauck, N., Lipovich, L., et al. (2000). Mapping and Genome Organization of Microsatellite Sequences in Rice (*Oryza sativa* L.). *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 100(5), 697-712.
- Templeton, A. R., Boerwinkle, E., and Sing, C. F. (1987). A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping. I. Basic theory and an analysis of alcohol dehydrogenase activity in *Drosophila*. *Journal of Genetics: The Genetics Society of America*, 117(2), 343-351.
- Thakur, M. C., Agrawal, S., Patel, M., Khan, A., and Doshi, H. V. (2013). Genetic Variability of *Nicotiana tabacum* (Linn.) Using SSR Marker. *World Journal of Agricultural Research*, 1, 124-129.
- Thi, H., Vu, T., Le, D. D., Ismail, A. M., and Le, H. H. (2012). *Marker-assisted backcrossing (MABC) for improved salinity tolerance in rice (Oryza sativa L.) to cope with climate change in Vietnam* Department of Molecular Biology, Agricultural Genetics Institute (AGI), Pham Van Dong Street, . Phillipine: International Rice Research Institute (IRRI)o. Document Number)
- Thoday, J. M. (1961). Location of polygenes. *Nature*, 191(1), 368-370.
- Thomson, M. J., de Ocampo, M., Egdane, J., Rahman, M. A., Sajise, A. G., Adorada, D. L., et al. (2010). Characterizing the Saltol quantitative trait locus for salinity tolerance in rice. *Rice*, 3(2-3), 148-160.

- Trethowan, R. M., and Mujeeb-Kazi, A. (2008). Novel germplasm resources for improving environmental stress tolerance of hexaploid wheat. *Crop Science*, 48(4), 1255-1265.
- Tsai, T. M., and Huang, H. J. (2006). Effects of iron excess on cell viability and mitogen-activated protein kinase activation in rice roots. *Physiologia Plantarum*, 127(4), 583-592.
- Tuong, T. P. (2000). Productive Water Use in Rice Production: Opportunities and Limitations. *Journal of Crop Production*, 2(2), 241-264.
- Ueno, D., NaokiYamaji, IzumiKono, FengHuang, C., Ando, T., Yano, M., et al. (2010). Gene limiting cadmium accumulation in rice. *Proceedings of the National Academy of Sciences*, 107(38), 16500-16505.
- Utz, H. F., and Melchinger, A. E. (1996). PLABQTL: A Program for Composite Interval Mapping of QTL. *Journal of Quant Trait Loci by U.S. Department of Agriculture*, 2(1), 1-5.
- van Beek, C. L., Meerburg, B. G., Schils, R. L., Verhagen, J., and Kuikman, P. J. (2010). Feeding the world's increasing population while limiting climate change impacts: linking N₂O and CH₄ emissions from agriculture to population growth. *environmental science & policy*, 13(2), 89-96.
- Violante, A., Cozzolino, V., Perelomov, L., Caporale, A. G., and Pigna, M. (2010). Mobility and Bioavailability of Heavy Metals and Metalloids in Soil Environments. *Journal of soil science and plant nutrition*, 10(3), 268-292.
- Vreugdenhil, D., Aarts, M. G. M., Koornneef, M., Nelissen, H., and Ernst, W. H. O. (2004). Natural variation and QTL analysis for cationic mineral content in seeds of *Arabidopsis thaliana*. *Journal of Plant, Cell & Environment*, 27(7), 828-839.
- Wah, C. A. (1998). *Direct Seeded Rice in Malaysia* (Vol. 1). FAO Regional Office for Asia and The Pacific, Bangkok: Asia-Pacific Association of Agricultural Research Institutions (APAARI) Publication.
- Wallace, J. G., Larsson, S. J., and Buckler, E. S. (2014). Entering the second century of maize quantitative genetics. *Heredity (Nature): Special Issue on Quantitative Genetics*, 112(1), 30-38.
- Wan, J.-L., Zhai, H.-Q., and Wan, J.-M. (2005). Mapping of QTLs for ferrous iron toxicity tolerance in rice (*Oryza sativa* L.). *Journal of genetics and genomics (Yi Chuan Xue Bao)*, 32(11), 1156-1166.

- Wan, J.-l., Zhai, H.-q., Wan, J.-m., and Ikehashi, H. (2003). Detection and analysis of QTLs for ferrous iron toxicity tolerance in rice, *Oryza sativa* L. *Euphytica*, 131(2), 201-206.
- Wang, D. L., Zhu, J., Li, Z. K., and Paterson, A. H. (1999). *User manual for QTLMapper version 1.0*. Unpublished manuscript, Texas A&M
- Wang, S. C., Basten, C. J., Zeng, Z. B., and Cartographer, W. Q. (2006). WinQtlCart V2.5. *Program in Statistical Genetics, North Carolina State University, Raleigh, NC*.
- Wang, Y.-H., Behera, T. K., and Kole, C. (2011a). *Genetics, Genomics and Breeding of Cucurbits (13th Edition)* (13 ed.): Taylor & Francis Group New York.
- Wang, Z., Chen, Z., Cheng, J., Lai, Y., Wang, J., Bao, Y., et al. (2012). QTL Analysis of Na⁺ and K⁺ Concentrations in Roots and Shoots under Different Levels of NaCl Stress in Rice (*Oryza sativa* L.). *PLoS ONE*, 7, 1-9.
- Wang, Z., Wang, J., Bao, Y., Wu, Y., and Zhang, H. (2011b). Quantitative trait loci controlling rice seed germination under salt stress. *Euphytica: International Journal of Plant Breeding*, 178(3), 297-307.
- Watanabe, T., and Osaki, M. (2001). Influence of aluminum and phosphorus on growth and xylem sap composition in *Melastoma malabathricum* L. *Plant and Soil: International Journal on Plant-Soil Relationships*, 237(1), 63-70.
- White, P. (2012). *Ion Uptake Mechanisms of Individual Cells and Roots: Short-distance Transport* (3 ed.). United States of America (USA): Elsevier.
- Williams, P. N., Price, A. H., Raab, A., Hossain, S. A., Feldmann, J., and Meharg, A. A. (2005). Variation in arsenic speciation and concentration in *paddy* rice related to dietary exposure. *Environmental Science & Technology*, 39(15), 5531-5540.
- Wissuwa, M., Yano, M., and Ae, N. (1998). Mapping of QTLs for phosphorus-deficiency tolerance in rice (*Oryza sativa* L.). *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 97(5-6), 777-783.
- Wu, F.-B., Jing, D., Jia, G.-X., Zheng, S.-J., and Zhang, G.-P. (2006). Genotypic difference in the responses of seedling growth and Cd toxicity in rice (*Oryza sativa* L.). *Agricultural Sciences in China*, 5(1), 68-76.

- Wu, L.-B., Shhadi, M. Y., Gregorio, G., Matthus, E., Becker, M., and Frei, M. (2014a). Genetic and physiological analysis of tolerance to acute iron toxicity in rice. *Rice*, 7(1), 1.
- Wu, L.-B., Shhadi, M. Y., Gregorio, G., Matthus, E., Becker, M., and Frei, M. (2014b). Genetic and physiological analysis of tolerance to acute iron toxicity in rice. *Rice*, 7(1), 8.
- Wu, P., Hu, B., Liao, C. Y., Zhu, J. M., Wu, Y. R., Senadhira, D., et al. (1998). Characterization of tissue tolerance to iron by molecular markers in different lines of rice. *Plant and Soil: International Journal on Plant-Soil Relationships*, 203(2), 217-226.
- Wu, P., Liao, C. Y., Hu, B., Yi, K. K., Jin, W. Z., Ni, J. J., et al. (2000). QTLs and epistasis for aluminum tolerance in rice (*Oryza sativa* L.) at different seedling stages. *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 100(8), 1295-1303.
- Wu, P., Luo, A., Zhu, J., Yang, J., Huang, N., and Senadhira, D. (1997). Molecular markers linked to genes underlying seedling tolerance for ferrous iron toxicity. In *Plant Nutrition for Sustainable Food Production and Environment* (pp. 789-792): Springer.
- Wu, Y.-P., Ko, P.-Y., Lee, W.-C., Wei, F.-J., Kuo, S.-C., Ho, S.-W., et al. (2010). Comparative analyses of linkage maps and segregation distortion of two F₂ populations derived from *japonica* crossed with *indica* rice. *Hereditas (Genetics & Heredity)*, 147(5), 225-236.
- Xiao, J., Li, J., Yuan, L., McCouch, S. R., and Tanksley, S. D. (1996). Genetic diversity and its relationship to hybrid performance and heterosis in rice as revealed by PCR-based markers. *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 92(6), 637-643.
- Xiong, Z. T., and Wang, H. (2005). Copper toxicity and bioaccumulation in Chinese cabbage (*Brassica pekinensis* Rupr.). *Journal of Environmental Toxicology*, 20(2), 188-194.
- Xu, K., Xu, X., Fukao, T., Canlas, P., Maghirang-Rodriguez, R., Heuer, S., et al. (2006). *Sub1A* is an ethylene-response-factor-like gene that confers submergence tolerance to rice. *nature*, 442(7103), 705-708.
- Xu, Y., Zhu, L., Xiao, J., Huang, N., and McCouch, S. R. (1997). Chromosomal regions associated with segregation distortion of molecular markers in F₂,

- backcross, doubled haploid, and recombinant inbred populations in rice (*Oryza sativa* L.). *Molecular and General Genetics MGG*, 253(5), 535-545.
- Xue, D., Chen, M., and Zhang, G. (2009). Mapping of QTLs associated with cadmium tolerance and accumulation during seedling stage in rice (*Oryza sativa* L.). *Euphytica*, 165(3), 587-596.
- Xue, Y., Wan, J. M., Jiang, L., Liu, L. L., Su, N., Zhai, H. Q., et al. (2006). QTL analysis of aluminum resistance in rice (*Oryza sativa* L.). *Plant and Soil: International Journal on Plant-Soil Relationships*, 287(1-2), 375-383.
- Yadav, S. K. (2010). Heavy metals toxicity in plants: an overview on the role of glutathione and phytochelatins in heavy metal stress tolerance of plants. *South African Journal of Botany*, 76(2), 167-179.
- Yao, M.-Z., Wang, J.-F., Chen, H.-Y., Zhai, H.-Q., and Zhang, H.-S. (2005). Inheritance and QTL mapping of salt tolerance in rice. *Rice: Applied Plant Science Journal*, 12(1), 25-32.
- Younas, M., Shahzad, F., Afzal, S., Khan, M. I., and Ali, K. (1998). Assessment of Cd, Ni, Cu, and Pb pollution in Lahore, Pakistan. *Environment International Journal*, 24(7), 761-766.
- Young, N. D. (2000). 3 . Constructing a plant genetic linkage map with DNA markers. 31-47.
- Young, N. D. (2001). Constructing a plant genetic linkage map with DNA markers. *DNA-Based Markers in Plants*, 31-47.
- Zhang, J., Zhu, Y.-g., Qian, Q., and Duan, G.-l. (2005a). Mapping quantitative trait loci associated with arsenic accumulation in rice (*Oryza sativa*). 2004-2009.
- Zhang, J. J., Lu, Y. C., Zhang, J. J., Tan, L. R., and Yang, H. (2014). Accumulation and toxicological response of atrazine in rice crops. *Ecotoxicology and environmental safety*, 102, 105-112.
- Zhang, W. P., Shen, X. Y., Wu, P., Hu, B., and Liao, C. Y. (2001). QTLs and epistasis for seminal root length under a different water supply in rice (*Oryza sativa* L.). *Theoretical and Applied Genetics: International Journal of Plant Breeding Research*, 103(1), 118-123.
- Zhang, X., Zhang, F., and Mao, D. (1998). Effect of iron plaque outside roots on nutrient uptake by rice (*Oryza sativa* L.). Zinc uptake by Fe-deficient rice. *Plant and Soil: International Journal on Plant-Soil Relationships*, 202(1), 33-39.

- Zhang, Y., Luo, L., Liu, T., Xu, C., and Xing, Y. (2009). Four rice QTL controlling number of spikelets per panicle expressed the characteristics of single Mendelian gene in near isogenic backgrounds. *Theoretical and applied genetics*, 118(6), 1035-1044.
- Zhang, Z.-H., Qu, X.-S., Wan, S., Chen, L.-H., and Zhu, Y.-G. (2005b). Comparison of QTL controlling seedling vigour under different temperature conditions using recombinant inbred lines in rice (*Oryza sativa* L.). *Annals of botany*, 95(3), 423-429.
- Zhou, G., Delhaize, E., Zhou, M., and Ryan, P. R. (2011). Biotechnological solutions for enhancing the aluminium resistance of crop plants. *Abiotic stress in plants—mechanisms and adaptations. Brisbane: InTech*, p119-142.
- Zhu, R., Gao, Y., and Zhang, Q. (2014). Quantitative trait locus mapping of floral and related traits using an F₂ population of *Aquilegia*. *Plant breeding*, 133(1), 153-161.
- Zong, G., Wang, A., Wang, L., Liang, G., Gu, M., Sang, T., et al. (2012). A Pyramid Breeding of Eight Grain-yield Related Quantitative Trait Loci Based on Marker-assistant and Phenotype Selection in Rice (*Oryza sativa* L.). *Journal of Genetics and Genomics*, 39(1), 335-350.